CUMULATIVE AUTHOR INDEX
Journal of Economic Entomology
Volume 105, 2012

A

Abney, M., 943
Abreu, A. G., 2136
Adamczyk, Jr., J. J., 1268
Adamson, D., 1115
Adgaba, N., 777
Adkins, S., 1447
Aguirre-Rojas, L., 2180
Ahn, Y.-J., 1329
Aikins, M. J., 282, 1107
Al Attal, Y. Z., 777
Al-Deeb, M. A., 2062
Alemandri, V., 48
Aleksandra, D., 670
Alfaress, S., 1629
Al-Ghamdi, A. A., 777
All, J., 1825
Allison, J. D., 923
Alma, A., 1311
Almqvist, C., 916
Aluja, M., 823
Alvarez, J. M., 783
Amrine, J. W., 1801
An, Z., 1915
Anciso, J. R., 1268
Anderson, D. J., 896
Anderson, J. F., 1789
Anderson, K. M., 1988
Anderson, P. F., 259
Anderson, T. D., 207
Andrić, G. G., 670
Angeli, G., 322
Ansari, M. J., 777
Appel, A. G., 1385
Argüello Caro, E. B., 48
Armstrong, J. S., 896
Arthur, F. H., 718
Arthurs, S. P., 1573
Ashley, R. O., 1988
Athanassiou, C. G., 282, 1088
Atiama-Nurbel, T., 1358
Atkinson, T. H., 533
Averill, A. L., 1366
Awondo, S. N., 943
Aydin, L., 1890

B

Babin, R., 1285
Bacheler, J. S., 1612
Backus, E. A., 1465, 1492
Bailey, W. C., 259, 1248
Baldessari, M., 322
Baluch, S. D., 642
Ban, L., 2129
Bansal, R., 1432
Bao, H., 739
Barr, N. B., 2147
Barrantes, L. D., 1350
Bartlett, J. S., 979
Basit, M., 1401
Baum, J. A., 616
Bedoukian, R., 1746
Beede, R. H., 1681
Beer, M.S.D., 1954
Beers, E. H., 1342, 1350
Behrens, N. S., 890
Bejerman, N., 48
Belton, P., 72
Bennett, G. W., 1736
Berenbaum, M. R., 410, 878
Bernal, J. S., 1532
Bester, G. G., 1268
Bethke, J. A., 753
Beuzelin, J. M., 1998
Bezirganoglu, I., 1581
Bharadwaj, A., 222
Bielza, P., 1006
Bilong, C.F.B., 1285
Blackshaw, R. E., 1261
Blais, M., 1929
Blanco, C. A., 2136
Blanco, M. H., 1457
Blount, N. A., 1640
Bommarco, R., 1620
Boquel, S., 1903
Borden, J. H., 304, 1668
Borges, P.A.V., 2213
Bostanian, N. J., 866
Boulogne, I., 1224
Bourgeois, A. L., 313
Bouyer, J., 1477
Boyd, E. A., 1335
Bradshaw, J. D., 1268
Brandenburg, R. L., 2045
Bravo, B., 1940
Breece, C. R., 1134
Breed, M. D., 1145
Brewer, M. J., 896
Briano, J. A., 817
Brijwani, M., 703, 709
Brockerhoff, E. G., 1694
Brown, A. E., 120
Brown, K. S., 518
Brown, R. S., 616
Bruck, D. J., 169, 1350
Bryant, J. L., 957
Buchwald, R., 1145
Buczkowski, G., 1736
Buntin, G. D., 1825
Buonaccorsi, J. P., 1366
Burd, J. D., 1057
Burkness, E. C., 1773
Burrack, H. J., 1350
Buschman, L. L., 598
Buss, E. A., 1573
Butler, C. D., 1509
Butler, R. C., 1694
Bynum, E. D., 1268
Byrne, F. J., 505, 753

C

Cabello, T., 2085
Cai, Y., 1503
Cakmak, I., 1890
Calcaterra, L. A., 817
Cameron, E. C., 1051
Campbell, D. M., 1694
Campbell, J. F., 686
Campbell, M., 304
Cang, T., 92
Cao, J., 583
Caragea, P., 259
Caren, J. R., 20
Carey, J. R., 371
Caro-Corrales, J., 1940
Carpio, L. A., 1268
Carrera, S. G., 2180
Carrière, Y., 994
Castagnaro, A. P., 573
Castañera, P., 214
Castle, S. J., 67, 1392
Catchot, A., 2007
Cave, R. D., 1719
Cervantes, F. A., 783
Chabi-Olaye, A., 161
Chamarthi, S. K., 1069
Chamberlin, J. R., 753
Chang, S.-C., 1100
Chen, H., 739
Chen, J.-L., 854
Chen, L., 92
Chen, M., 354
Chen, Q.-Y., 1566
Chen, Z.-M., 1034
Cheng, D.-F., 854
Cheng, C., 1471
Cheng, X., 1164
Cheng, Z., 1164
Chernet, M. H., 777
Chi, H., 930
Chinta, S. P., 497
Chirino, M. G., 386
Chiroleu, F., 1358
Chon, I., 1540
Chu, D., 242, 930
Chynoweth, R., 1006
Cibils-Stewart, X., 1407

CUMULATIVE AUTHOR INDEX
Journal of Economic Entomology
Volume 105, 2012

A

Abney, M., 943
Abreu, A. G., 2136
Adamczyk, Jr., J. J., 1268
Adamson, D., 1115
Adgaba, N., 777
Adkins, S., 1447
Aguirre-Rojas, L., 2180
Ahn, Y.-J., 1329
Aikins, M. J., 282, 1107
Al Attal, Y. Z., 777
Al-Deeb, M. A., 2062
Alemandri, V., 48
Aleksandra, D., 670
Alfaress, S., 1629
Al-Ghamdi, A. A., 777
All, J., 1825
Allison, J. D., 923
Alma, A., 1311
Almqvist, C., 916
Aluja, M., 823
Alvarez, J. M., 783
Amrine, J. W., 1801
An, Z., 1915
Anciso, J. R., 1268
Anderson, D. J., 896
Anderson, J. F., 1789
Anderson, K. M., 1988
Anderson, P. F., 259
Anderson, T. D., 207
Andrić, G. G., 670
Angeli, G., 322
Ansari, M. J., 777
Appel, A. G., 1385
Argüello Caro, E. B., 48
Armstrong, J. S., 896
Arthur, F. H., 718
Arthurs, S. P., 1573
Ashley, R. O., 1988
Athanassiou, C. G., 282, 1088
Atiama-Nurbel, T., 1358
Atkinson, T. H., 533
Averill, A. L., 1366
Awondo, S. N., 943
Aydin, L., 1890

B

Babin, R., 1285
Bacheler, J. S., 1612
Backus, E. A., 1465, 1492
Bailey, W. C., 259, 1248
Baldessari, M., 322
Baluch, S. D., 642
Ban, L., 2129
Bansal, R., 1432
Bao, H., 739
Barr, N. B., 2147
Barrantes, L. D., 1350
Bartlett, J. S., 979
Basit, M., 1401
Baum, J. A., 616
Bedoukian, R., 1746
Beede, R. H., 1681
Beer, M.S.D., 1954
Beers, E. H., 1342, 1350
Behrens, N. S., 890
Bejerman, N., 48
Belton, P., 72
Bennett, G. W., 1736
Berenbaum, M. R., 410, 878
Bernal, J. S., 1532
Bester, G. G., 1268
Bethke, J. A., 753
Beuzelin, J. M., 1998
Bezirganoglu, I., 1581
Bharadwaj, A., 222
Bielza, P., 1006
Bilong, C.F.B., 1285
Blackshaw, R. E., 1261
Blais, M., 1929
Blanco, C. A., 2136
Blanco, M. H., 1457
Blount, N. A., 1640
Bommarco, R., 1620
Boquel, S., 1903
Borden, J. H., 304, 1668
Borges, P.A.V., 2213
Bostanian, N. J., 866
Boulogne, I., 1224
Bourgeois, A. L., 313
Bouyer, J., 1477
Boyd, E. A., 1335
Bradshaw, J. D., 1268
Brandenburg, R. L., 2045
Bravo, B., 1940
Breece, C. R., 1134
Breed, M. D., 1145
Brewer, M. J., 896
Briano, J. A., 817
Brijwani, M., 703, 709
Brockerhoff, E. G., 1694
Brown, A. E., 120
Brown, K. S., 518
Brown, R. S., 616
Bruck, D. J., 169, 1350
Bryant, J. L., 957
Buchwald, R., 1145
Buczkowski, G., 1736
Buntin, G. D., 1825
Buonaccorsi, J. P., 1366
Burd, J. D., 1057
Burkness, E. C., 1773
Burrack, H. J., 1350
Buschman, L. L., 598
Buss, E. A., 1573
Butler, C. D., 1509
Butler, R. C., 1694
Bynum, E. D., 1268
Byrne, F. J., 505, 753

C

Cabello, T., 2085
Cai, Y., 1503
Cakmak, I., 1890
Calcaterra, L. A., 817
Cameron, E. C., 1051
Campbell, D. M., 1694
Campbell, J. F., 686
Campbell, M., 304
Cang, T., 92
Cao, J., 583
Caragea, P., 259
Caren, J. R., 20
Carey, J. R., 371
Caro-Corrales, J., 1940
Carpio, L. A., 1268
Carrera, S. G., 2180
Carrière, Y., 994
Castagnaro, A. P., 573
Castañera, P., 214
Castle, S. J., 67, 1392
Catchot, A., 2007
Cave, R. D., 1719
Cervantes, F. A., 783
Chabi-Olaye, A., 161
Chamarthi, S. K., 1069
Chamberlin, J. R., 753
Chang, S.-C., 1100
Chen, H., 739
Chen, J.-L., 854
Chen, L., 92
Chen, M., 354
Chen, Q.-Y., 1566
Chen, Z.-M., 1034
Cheng, D.-F., 854
Cheng, C., 1471
Cheng, X., 1164
Cheng, Z., 1164
Chernet, M. H., 777
Chi, H., 930
Chinta, S. P., 497
Chirino, M. G., 386
Chiroleu, F., 1358
Chon, I., 1540
Chu, D., 242, 930
Chynoweth, R., 1006
Cibils-Stewart, X., 1407

Čičková, H., 62
Cifuentes, D., 1006
Cilas, C., 1285
Clark, T. L., 616
Cleaver, C. M., 1645
Cloyd, R. A., 505, 2097
Coats, J. R., 890
Coll, M., 1520
Collins, H. L., 354
Conder, N., 344
Cossé, A., 429
Costello, M. I., 1293
Cônsoli, F. L., 2136
Cook, D. R., 2007
Cook, M. A., 1205
Cook, P., 1557
Cooper, W. R., 860, 1057
Cooperband, M. F., 1694
Costa-Júnior, L. M., 1979
Coulin, C., 817
Cowell, B., 2076
Cowles, R. S., 1789
Craig, C., 1825
Crook, D. J., 429
Crosslin, J. M., 1268
Cui, F., 232
Cullen, E., 259
Cumsille, R., 1844
Czokajlo, D., 2076

D

Daane, K. M., 1681
Dabney, C. P., 149, 1668
Dahlgren, L., 1895
Daly, Z., 1130
Danka, R. G., 313
Dashiell, K. E., 608
Dastjerdi, H. R., 1439
Davey, R. B., 2207
Davis, H., 2180
Davis, R. W., 1766
de Barro, P., 48
de Guzman, L. I., 313
de Kogel, W. J., 169
de la Poza, M., 214
De La Rúa, P., 1006
Deguine, J.-P., 1358
Delaney, D. A., 20
DeLucia, E. H., 878
Demark, J. J., 533
Dennehy, T. J., 753
DePalma, E., 1077
Desfontaines, L., 1224
Desneux, N., 1277
Deutschlander, M., 632
Deyrup, M. A., 659
Dhavala, S., 1825
Diaz-Montano, J., 632
Dibog, L., 1285
Dickey, A. M., 753
Diffie, S., 40, 1447
DiFonzo, C., 259
Dively, G. P., 207
Djieto-Lordon, C., 1285
Domingues, F. A., 2136
Dosdall, L. M., 1261
Doskocil, J. P., 2045
Dossett, M., 1465
Dou, W., 560
Douraguia, E., 1358
Dreves, A. J., 1350
Du Tott, C. L. N., 1963
Du, Y.-J., 1566
Duan, J. J., 792
Duan, L. Q., 344
Duell, M., 1890
Dumón, A. D., 48
Duso, C., 176
Duyck, P.-F., 371

E

Echegaray, E. R., 2097
Eger, J. E., 533
Ekasi, S., 1963
Ekesi, S., 2068
Ellersieck, M. R., 1248
Elliott, K., 259
Ellis, M. D., 1895
Endo, N., 1024
England, G. K., 480
Epsky, N. D., 471, 659
Erhardt, S., 1688
Esmaeili, N., 1809

F

Fail, J., 632
Fan, J., 1164, 1503
Farinś, G. P., 214
Faroni, L.R.A., 2187
Fathipour, Y., 2172
Fehr, W. R., 252
Feng, C.-H., 1566
Feng, D.-D., 1034
Feng, Hongqiang, 1921
Feng, Hongyun, 1921
Feng, X., 739
Fernandes, F. S., 113
Fernandez, F. J., 2085
Ferracini, C., 1311
Ferrari, E., 1311
Ferreira, M. T., 2213
Fettig, C. J., 149, 1668
Fields, P. G., 718
Fierke, M. K., 2029
Fitzpatrick, S. M., 1205
Flasinski, S., 616
Flinn, P. W., 703, 709
Florane, C., 1746
Folgarait, P. J., 386
Follett, P. A., 817, 1178, 1971
Fonsah, E. G., 943
Foottit, R., 1909
Fornasiero, D., 176
Fowler, D. L., 1668
Francese, J. A., 1929
Francis, F., 854
Frankenhuyzen, K. V., 1929
Fraser, I., 429
French, B. W., 1407
French, R., 1801
Friesen, K., 726
Fu, W.-J., 549
Furlong, M. J., 1115

G

Gaiotti, F., 176
Gallego, J. R., 2085
Gamez, M., 2085
Gan, Y., 1261
Ganbalani, G. N., 1439
Gao, Changsheng, 242
Gao, Congfen, 2129
Garcia, M. G., 573
Gardescu, S., 2015
Gassmann, A. J., 994, 1407
Gastaminza, G., 573
Gautam, B. K., 1766
Ge, Y. D., 1095
George, J. E., 2207
Germosén-Robineau, L., 1224
Giguére, M.-A., 1903, 1909
Gilbert, L. E., 386
Gilchrist, A. S., 1051
Gilrein, D., 753
Gimmel, M. L., 923
Gispert, C., 1392
Gitaitis, R. D., 40
Gitonga, L. M., 2068
Golizadeh, A., 592, 1439, 1809
Gómez-Torres, M. L., 338
Gondim, M.G.C., 1419
Gong, L., 232
Goodrich, B. A., 1645
Goolsby, J. A., 1268
Goosey, H. B., 102
Gore, J., 2007
Gordon, J. R., 971
Gotoh, T., 1043
Gould, F., 767
Graham, E. E., 395, 837
Gratton, C., 259
Green, T.R., 374
Greene, J., 2007
Griepink, F. C., 169
Groot, A. T., 573
Grové, T., 1954
Gu, P., 1566
Guedes, R.N.C., 2187
Guerra, I. M., 451
Guertin, C., 1929
Guillén, J., 1006
Guillén, L., 823
Gulzar, A., 964
Guo, W.-C., 549
Guo, X.-L., 1157
Guo, Y., 1915

H
Haga, A. V., 1878
Hajek, A. E., 2015
Haklova, M., 1487
Hall, D. G., 783
Hall, P. M., 753
Hama, M., 1781
Hamby, K. A., 1350
Hammon, R. W., 1057
Han, P., 1277
Hanks, L. M., 906, 2029
Hannon, E. R., 994
Hanson, B., 1988
Hanula, J. L., 461
Hardin, J. G., 1645
Hardman, J. M., 866
Haribalan, P., 1329
Harker, K. N., 1261
Harpaz, T. S., 1520
Harris, J. W., 313
Harris, M. O., 1988
Harrison, Jr., H. F., 651
Hartzer, M., 709
Hasan, M. M., 282, 1107
Hasanpour, M., 592
Hasegawa, E., 555
Hassan, E., 1379
Hauer, R. H., 196
Haviland, D. R., 1350, 1681
Hay-Roe, M.G.M.M., 418
He, Y., 2129
He, Y.-R., 1157
Heath, R. R., 471, 659
Heck, G. R., 616
Hein, G. L., 1801
Hemati, S. A., 1439
Henderson, G., 1758, 1766
Hendrix, W., 1825
Henne, D. C., 1268
Herbert, D. A., 884, 1612
Herms, D. A., 1
Hernández, E., 1940
Hernandez-Suarez, E., 2085
Hesler, L. S., 608
Hesler, S., 936
Heather, P. N., 1379
Hibbard, B. E., 1248
Higbee, B. S., 1702
Higgins, R. A., 598
Hillbur, Y., 1988
Hinomoto, N., 1043
Hirase, K., 1781
Hitchner, E. M., 207
Hoddle, D., 402
Hoddle, H. S., 402
Hoddle, M., 1077
Hoffmann, E. J., 67
Hofstetter, R. W., 2107
Hollingsworth, R. G., 1238
Hong, X.-Y., 2161
Howard, R. H., 777
Hranitz, J. M., 1890
Htwe, A. N., 802
Hu, X., 242
Hu, X. P., 2115
Hu, Z., 739
Huang, D., 288
Huang, R.-N., 1591
Huang, T.-I, 847
Huang, Z., 2129
Huber, D.P.W., 149, 1668
Hulasare, R., 709
Hulbert, D., 1688
Husseneder, C., 518
Hutchison, W. D., 1773
Hwang, S.-Y., 1581

I
Ichiki, R. T., 802
Ingegno, B. L., 1311
Isaacs, R., 936, 1350, 1688

J
Jackson, D. M., 651
Jackson, R., 2007
Jacobi, W. R., 1645
Jacome, I., 1548
Jamieson, L. E., 1171
Jang, E. B., 54
Jayas, D. S., 1194
Jeske, D. R., 1077
Jian, F., 1194
Jiang, H.-B., 560
Jiang, W.-H., 549
Jiao, Z., 1603
Jin, Q.-A., 988
Jin, T., 988
Jin, Y., 1921
Johnson, D., 259
Johnson, D. T., 2076, 2196
Johnson, E. N., 1261
Johnson, R. M., 1895
Johnson, W. A., 1629
Johny, S., 1929
Jones, G. C., 1929
Jones, M. M., 1426
Jones, R. W., 1871
Jones, S. C., 957
Joseph, S. V., 1302
Joshi, A., 1268
Joubert, P. H., 1954
Juárez, M. L., 418, 573

K
Kachigamba, D. L., 2068
Kafle, L., 1095
Kalischuk-Tymensen, L. D., 1863
Kang, L., 232
Karimi, S., 2172
Kasai, S., 1796
Kawahara, N., 1781
Keddie, B. A., 1878
Keller, J. J., 20
Kemp, K. E., 505
Kemp, W. P., 14
Kendra, P. E., 659
Khrimian, A., 429
Kim, J.-R., 1329
Kim, S., 1524
Kim, S.-H.S., 2076
Kim, Y., 1540
Kindlmann, P., 1487
Kitchens, S. C., 1640
Klinkhamer, P.G.L., 1816
Kljajić P. J., 670
Kobayashi, K., 555
Kobayashi, M., 1796
Koch, F. H., 438
Komagata, O., 1796
Kong, J., 288
Konopka, J. K., 872
Kouloussis, N. A., 371
Kovacs, E., 304
Kozánek, M., 62
Krüger, K., 161
Krupke, C. H., 259
Kuhar, T., 1612
Kuhar, T. P., 207
Kuo, T.-C., 1591
Kurashima, R., 54, 1186
Kutcher, H. R., 1261
Kyei poku, G., 1929

L
Labruna, M. B., 1979
Lacroix, M., 1238
Lagos, D., 259
Lai, L.-C., 1591
Lai, X.-F., 1034
Lait, C. G., 304, 1668
Lamb, B. K., 451
Lambkin, T. A., 979
Langemeier, M. R., 709
Lara, J. R., 1077
Larson, R., 1194
Lavallée, R., 1929
Lawrance, A., 410
Lax, A., 1746
Lee, C.-Y., 1100
Lee, D.-H., 1726
Lee, J. C., 1350, 1465
Lee, L.-C., 533
Lees, M. D., 533
Lefebvre, M., 866
Leiss, K. A., 1816
Leonard, B. R., 1825, 2007
Leskey, T. C., 1548, 1726
Lester, P. J., 1139
Lewis, B. A., 2076
Lewis, P. A., 2015
Li, G.-Q., 549
Li, Guanghong, 1503
Li, Guoping, 1921
Li, H.-S., 2161
Li, Jie, 583
Li, Juan, 1566

Li, P., 1149
Li, X., 242
Li, Z., 739
Liburd, O. E., 480
Liedo, P., 1234
Lightle, D. M., 1465
Lim, U. T., 1524
Lima, D. B., 1419
Lin, Q., 739
Lin, Y.-Y., 988
Lin, Z., 232
Little, N. S., 1640
Liu, T.-X., 490
Liu, X., 354, 583
Liu, Y., 854
Liu, Y.-B., 810
Loeb, G., 936
Lohmeyer, K. H., 2207
Londo, A. J., 1640
Lopes, A.J.O., 1979
Lopes, J.R.S., 329
Loranger-Merciris, G., 1224
Lowenberger, C. A., 72
Lu, K.-H., 1100
Lu, W.-P., 549
Lu, Y., 1603
Lu, Z., 2122
Ludwig, S., 753
Lundin, O., 1620
Lundquist, J. E., 1659
Luo, H., 1566
Luo, Y.-L., 1157
Luthe, D. S., 120
Lyon, C. A., 1145
Lyons, D. B., 1929
Lysyk, T. J., 732, 1863
Lytle, J. M., 1532

M

Ma, H.-J., 1034
Machial, C. M., 304
Mackey, B., 1557
Magarey, R. D., 438
Mahmoud, S. T., 2062
Mahob, R., 1285
Mainali, B. P., 1524
Malaquias, J. B., 113
Malone, S., 884, 1612
Mamidala, P., 1432
Mangan, R. L., 363
Marijana P. G., 670
Marques, R. N., 329
Mascarin, G. M., 1573
Mason, C. E., 1214
Mason, K. S., 936
Mastro, V. C., 429
Mathieson, M., 1139
Matsuda, T., 1043
Matsukura, K., 129
Matsumura, M., 129
Mattio, M. F., 48
Mauffette, Y., 866
Maw, E., 1909
Mayfield, III, A. E., 461
Mazzoni, V., 322
Mbata, G. N., 679
McCarville, M. T., 1835
McConnell, T. E., 1640
McCornack, B. P., 259, 1629
McCullough, D. G., 1, 272, 837
McElfresh, J. S., 497, 2029
McInnis, D. O., 54
McKelvey, S. R., 149, 1668
McKenney, J. L., 923
McKenzie, C. L., 753
McNeil, J. N., 872
McPheron, B. A., 2147
Meagher, R. L., 418, 1573
Meats, A. W., 1051
Meikle, W. G., 26
Melhim, A., 1130
Ming, Q.-L., 1471
Méndez, T., 1844
Mercader, R. J., 272
Messenger, M. T., 533
Meyer, S. E., 616
Mian, M.A.R., 1432
Michel, A. P., 1432
Midgarden, D., 471
Milan A., 670
Millar, J. G., 497, 837, 906, 949, 2029, 2035, 2052
Miller, D. R., 923
Miller, J. A., 2207
Miresmailli, S., 878
Mirjana M. M., 670
Mirnezhad, M., 1816
Mitchell, R. F., 906
Mittapalli, O., 1432
Mochizuki, A., 802
Moeri, O. E., 304
Mohamadi, P., 592
Mohamed, S. A., 1963
Monteiro, V. B., 1419
Montemayor, C. O., 1719
Montgomery, W. S., 659
Montoya, P., 1940
Morávek, I., 62
Moreno, A. T., 363
Morse, J. G., 497, 1532, 1557, 2035
Mosser, L. K., 1694
Mosti, M., 1311
Munyaneza, J. E., 1268
Murata, M., 802
Murúa, M. G., 573
Murugan, M., 2180
Musser, F., 2007
Musser, F. R., 1825
Musundire, R., 161
Muza, A., 936

N

Nabity, P. D., 878
Nagoshi, R. N., 418
Nakamura, S., 802
Nakao, T., 1781
Nanayakkara, U. N., 1903, 1909
Naoi, A., 1781
Narasu, L. M., 1069
Nascimento, A.R.B., 113
Nascimento Júnior, J. L., 113
Nascimento Júnior, J.R.S., 1979
Naseri, B., 592, 1439, 2172
Nault, B. A., 632
Nava, D. E., 338
Navone, P., 1311
Nay, J. E., 1335
Nderitu, P. W., 1963
Ndung'u, M. W., 2068
Neese, P. A., 533
Neven, L. G., 297, 1540
Nguyen, H. T., 802
Ni, X., 1457
Nichols, R. L., 242
Nicholson, R. A., 72
Nie, X., 1903, 1909
Niide, T., 598
Niogret, J., 659
Nishimoto, J., 1186
Niu, C.-Y., 1277
Niu, G., 410
Niu, J., 583
Nolting, S. P., 1825

O

O'Donovan, J. T., 1261
Oetting, R. D., 753
Ohm, H. W., 642
Omoto, C., 2136
O'Neal, M. E., 252, 259, 1835
O'Neil, R., 259
Onstad, D., 354
Ontivero, M., 573
Opit, G. P., 1107
Oppel, C., 792
Orbán, L. L., 34
Ordano, M., 823
Orman, A., 1890
Orpet, R., 878
Ortego, F., 214
Ortiz-Martínez, S. A., 1844
Oruc, H. H., 1890
Osborne, L. S., 753
Ottea, J., 971
Otvos, I. S., 344
Owen, C. L., 2147
Owens, C. B., 518
Ozier-Lafontaine, H., 1224

P

Pang, Y., 1164, 1503
Pant, H., 135
Papadopoulos, N. T., 371

Pappu, H. R., 40
Park, C. G., 379
Park, K. C., 379
Parman, V., 523
Parra, J.R.P., 338
Pascual-Villalobos, M. J., 679
Patt, J. M., 26
Pavan, F., 176
Payton, M. E., 679
Peairs, F. B., 1057
Peck, S. L., 1477
Pelletier, Y., 1903, 1909
Peng, C.-L., 1566
Peng, L.-N., 490
Peng, Y., 2129
Peng, Z.-Q., 988
Penn, S. R., 616
Perera, C. O., 1171
Perera, O. P., 2136
Pernal, S. F., 1878
Peterson, E. M., 1335
Peterson, R.K.D., 85
Petzold-Maxwell, J. L., 1407
Philips, C., 1612
Philips, C. R., 207
Phillips, T. W., 282, 1107
Picard, I., 1238
Pickett, B., 964
Piñero, J. C., 1548
Pino, M. D., 2085
Ploetz, R. C., 659
Plowright, C.M.S., 34
Plowright, R. C., 34
Poland, T. M., 395, 837
Pollock, H. S., 410
Popa, R., 374
Porter, P., 1268
Potter, D. A., 1320
Pound, J. M., 2207
Pozzebon, A., 176
Prabhaker, N., 1392
Prado, A. P., 1979
Prischmann, D. A., 608
Prokopy, R. J., 1548
Pruett, G. E., 659
Pszczolkowski, M. A., 2076
Purwatiningsih,, 1379
Pureswaran, D. S., 140
Puterka, G. J., 1057

Q

Qin, Y., 1149
Quek, S.-Y., 1171
Quilici, S., 371, 1358

R

Racette, G., 866
Radloff, S. E., 777
Ragenovich, I. R., 451
Ragsdale, D. W., 259
Raina, A., 1746
Ram C., 1844
Ramalho, F. S., 113
Randolph, T. L., 1057
Rascoe, J., 2052
Ravuiwasa, K. T., 1581
Razmjou, J., 592
Razza, J. M., 1214
Reagan, T. E., 1597, 1998
Reber, J., 1988
Redmond, C. T., 1320
Reeb, P., 1688
Reeck, G., 232
Regmi, H., 1095
Reich, R. M., 1659
Reisig, D. D., 884, 1612
Rendón, P., 471
Reut, M., 2076
Rhodes, E. M., 480
Rice, E. S., 1751
Rice, S. J., 979
Richardson, T. A., 1350
Riegel, C., 518
Riggins, J. J., 1640
Riley, D. G., 40, 943, 1302, 1447
Rimoldi, F., 1982
Rinderer, T. E., 313
Rinehart, J. P., 14
Rivera, P., 1940
Robertson, J. L., 1426
Robinson, L. J., 497, 2035
Roda, A., 2052
Rodriguez, S. M., 48
Roge¯s, M. E., 540, 1492
Roh, H. S., 379
Roitberg, B. D., 1205
Rojas, P., 490
Ronco, A. E., 1982
Rosenberg, O., 916
Ross, D. W., 451
Roush, R., 354
Rucker, A., 1712
Ruiz arce, R., 2147
Rull, J., 823
Rundlöf, M., 1620
Rush, C. M., 1268
Ryan-Bohac, J. R., 651
Ryba, S., 1487

S

Saeed, S., 1401
Sagili, R. R., 1134
Saleem, M. A., 1401
Salifu, D., 161
Salmieri, S., 1238
Sammataro, D., 26
Samuel, L., 1825
Sánchez, D., 1234
Sanchez, J. S., 659
Sandoya, G. V., 120
Sasaki, T., 129
Saunders, M., 936
Sayyed, A. H., 964, 1401
Schapaugh, W. T., 598
Scheffrahn, R. H., 2213
Scherer, C. W., 1736
Schidlo, N., 1816
Schleier, III, J. J., 85
Schmale, L., 753
Schmidt, N. P., 259
Schneider, M. I., 1982
Schuettpelz, N. M., 196
Schultz, T. P., 1640
Scott, I. M., 872
Selinger, L. B., 732, 1863
Semeao, A. A., 686
Seng, Y. E., 288
Serikawa, R. H., 1492
Shabbir, A., 1115
Sharif, E. M., 2062
Sharma, H. C., 1069
Shatters, Jr., R. G., 753
Shearer, P. W., 1350, 1712
Shelly, T., 1186
Shelton, A. M., 354, 632
Sheng, C., 232
Shi, W., 1915
Shi, X., 616
Shiba, T., 129
Shih, C. J., 1095
Shik, J. Z., 1751
Shivaji, R., 120
Short, B. D., 1726
Showler, A. T., 1597, 1998
Shukle, J. T., 642
Shu-Sheng, L., 1115
Siebert, M. W., 1825
Siegel, J. P., 410, 1702
Siegert, N. W., 272
Siegfried, B. D., 1895
Sievert, K., 726
Silva, C.A.D., 113
Silva, C. G., 1979
Silva, G. N., 2187
Silva, H.N.D., 1171
Silva, R., 1115
Silva-Brandão, K. L., 2136
Silverman, J., 1751
Sims, S. R., 1385
Singh, N., 2196
Singh, R. N., 1043
Siqueira, H.A.A., 1419
Siriwetwiwat, B., 1801
Sisterson, M. S., 994
Skabeikis, D. D., 2029
Sloderbeck, P. E., 1268
Sloderbeck, Phillip E., 686
Smith, C. L., 120
Smith, C. M., 598, 2180
Smith, H. G., 1620
Smith, W. D., 438
Smith, W.E.C., 120
Snodgrass, G. L., 2007
Snook, K., 1971
Solórzano, E.D.J., 1234
Son, B.-K., 1329

Son, Y., 1540
Sorucu, A., 1890
Souder, S. K., 1557
Sousa, A. H., 2187
Sparks, A., 1447
Sparks, D. L., 120
Spurgeon, D. W., 860
Srinivasan, R., 40, 783, 1302, 1447
Stafford, III, K. C., 222
Stanley, C. A., 1350
Stark, J. D., 1557
Stelinski, L. L., 540
Stewart, S. D., 1825
Stock, S. P., 994
Stocks, I., 2052
Stopka, P., 1487
Strand, T. M., 451
Striman, B. L., 906
Stringer, L. D., 1694
Su, N.-Y., 518
Subbarao, S., 616
Subramanyam, B., 703, 709
Suckling, D. M., 1, 54, 1694
Sugeno, W., 802
Sukuru, U. R., 616
Sullivan, B. T., 140
Sun, J.-R., 854
Sun, S., 1164, 1503
Sundaraj, S., 40
Supak, J. R., 1268
Sutherland, A., 1145
Sved, J. A., 1051
Swain, A. J., 979
Sylvester, H. A., 313

T

Tabashnik, B. E., 767, 994
Takáč, P., 62
Takakura, K. I., 1856
Takano, S.-I., 802
Takeychi, H., 1024
Talebi, A. A., 2172
Tan, C.-W., 1581
Tang, P.-A., 560
Tao, Y. L., 930
Tarpy, D. R., 20
Tavella, L., 1311
Taylor, D. B., 726
Teal, P.E.A., 2068
Teale, S. A., 2029
Tedeschi, R., 322
Teixeira, D. C., 329
Teixeira, L.A.F., 936
Tewari, S., 1366
Thistle, H. W., 451
Thistlewood, H. A., 866
Thomas, D. B., 2147
Thoms, E. M., 533
Throne, J. E., 282
Tilmon, K. J., 259
Timer, J., 936
Titera, D., 1487
Tiwari, S., 540
Tobin, P. C., 1
Toews, M. D., 847
Tofangsazi, N., 1573
Toft, R., 1139
Toledo, J., 1940
Tollerup, K. E., 1712
Tolley, M. P., 533
Tomasi, D., 176
Tomita, T., 1796
Torres-Ruiz, A., 1871
Torto, B., 2068
Tripathi, S., 135
Trona, F., 322
Troxclair, N. N., 1268
Trumble, J. T., 490, 1509
Truol, G., 48
Tuffly, M., 1659

U

Ugine, T. A., 2015

V

Vandame, R., 1234
Vandervoort, C., 1688
Vannatta, A. R., 196
van Tol, R.W.H.M., 169
Vargas, R. I., 1557
Vargo, E. L., 523, 533
Vassilakos, T. N., 1088
Vera, M. T., 573, 1178
Verdugo, J. A., 1844
Veysey, S. T., 1293
Vijay, P. M., 1069
Vila, E., 2085
Vilardi, J. C., 573
Villa, J. D., 313
Villagrán, M. E., 1178
Villanueva, R. T., 896
Villatoro, D., 471
Voegtlin, D., 259

W

Wagener, C. M., 313
Waliwitiya, R., 72
Walsh, D. B., 1350
Walton, V. M., 1350
Wang, B., 2167
Wang, C., 1758
Wang, D.-S., 1157
Wang, F., 1503
Wang, J.-J., 560
Wang, M., 739
Wang, Q., 92
Wang, X., 1019
Wang, Y., 92, 344
Ware, A. B., 1963
Waterworth, R. A., 949, 2035
Weersink, A., 1130
Weigle, T., 936
Weihman, S., 2052
Weinzierl, R. A., 1426
Weisz, R., 1612
Wen, H.-B., 988
Weslien, J., 916
White, N.D.G., 1194
Whitworth, J., 259
Whitworth, J. L., 783
Whitworth, R. J., 598, 686, 1629
Wiarda, S. L., 252
Wijayaratne, L.K.W., 718
Willett, F. J., 1268
Williams, C. E., 642
Williams, D. W., 1320
Williams, K. A., 505
Williams, W. P., 120
Willink, E., 418, 573, 1178
Wilson, B. E., 1597, 1998
Wilson, J. P., 1457
Wise, J. C., 1688
Wong, J.C.H., 906
Woods, B., 54
Woolf, A. B., 1171
Wright, D. J., 964
Wright, S. E., 1548, 1726
Wu, C., 92
Wu, K., 1603
Wu, T., 2167
Wu, W.-J., 1591
Wu, Y., 1019

X

Xia, Z.-H., 549
Xiao, L., 2167
Xie, H.-C., 854
Xu, H., 2122
Xu, J., 232
Xu, L. B., 344
Xu, M.-Y., 1034
Xu, W., 1457

Y

Yamamoto, P. T., 329
Yang, K., 1503
Yang, Y., 2122
Yao, M.-C., 1100
Yede, 1285
Yemshanov, D., 438
Yin, F., 739
Yocum, G. D., 14
Yokoyama, V. Y., 186
Youngman, R. R., 207
Yturralde, K. M., 2107
Yu, Q.-Y., 1034
Yu, R., 92
Yuan, Z., 2115

Z

Zalom, F. G., 295, 1350
Zalucki, M. P., 739, 979, 1115
Zechmann, B. J., 1268
Zens, B. A., 1268
Zhang, D., 739

Zhang, J., 1903, 2167
Zhang, Q., 354, 583
Zhang, S., 2129
Zhang, X.-F., 2161
Zhang, Y.-E., 1034
Zhang, Y.-M., 490
Zhang, Z., 1034
Zhao, D.-X., 2161
Zhao, H., 242
Zhao, X., 92
Zheng, X., 2122
Zhong, Y., 2167
Zhou, H.-B., 854
Zhou, X.-W., 1277
Zhu, J., 890, 1149
Zhu, J. J., 726
Zukoff, S. N., 1248
Zulhendri, F., 1171

CUMULATIVE SUBJECT INDEX
Journal of Economic Entomology
Volume 105, 2012

A

Abbott's formula,
  Bayesian statistics, 1856
  EC_{50}, 1856
  range estimation, 1856
  repellency, 1856
Acer platanoides,
  Anoplophora glabripennis, 2015
  fecundity, 2015
  longevity, 2015
  neonicotinoid, 2015
Aceria tosichella,
  Aceria tulipae, 1801
  wheat curl mite, 1801
  wheat streak mosaic, 1801
Aceria tulipae,
  Aceria tosichella, 1801
  wheat curl mite, 1801
  wheat streak mosaic, 1801
acetamiprid, 988, 1401
  avermectin, 988
  Brontispa longissimi, 988
  β-cypermethrin, 988
  fitness cost, 1401
  insecticide susceptibility, 988
  resistance, 1401
  whitefly, 1401
acetophenone, 149, 1668
  Dendroctonus ponderosae, 149
  nonhost angiosperm volatile, 149, 1668
  pest management, 149, 1668
  Pinus ponderosa, 149, 1668
  verbenone, 1668
acetylcholinesterase insensitivity,
  carbamate, 540
  general esterase, 540
  glutathione-S-transferase, 540
  huanglongbing, 540
acquisition,
  aphid behavior, 1903
  artificial diet, 1903
  potato virus Y, 1903
acrinathrin,
  cytochrome P450, 1006
  insecticide resistance, 1006
  over-expression, 1006
  western flower thrip, 1006
acute toxicity,
  Apis mellifera anatoliaca, 1890
  flumethrin, 1890
  oral, 1890
adaptation,
  Diuraphis noxia, 1057
  genetic diversity, 1057
  plant resistance, 1057
  wheat, 1057
adult longevity,
  emergence, 1524
  Gryon japonicum, 1524
  Ooencyrtus nezarae, 1524
  parasitism, 1524
adult mortality,
  Bacillus thuringiensis(*Bt*), 1863
  diet, 1863
  Stomoxys calcitrans, 1863
adult survival,
  Bactrocera oleae, 186
  humidity, 186
  reproductive biology, 186
  temperature, 186
Aethina tumida,
  diet, 26
  population growth, 26
  resource acquisition, 26
  temperature, 26
AFB,
  Apis mellifera, 1487
  honey bee, 1487
  matrix, 1487
  polymerase chain reaction (PCR), 1487
Africa,
  Glossinidae, 1477
  insect control, 1477
  livestock disease, 1477
  simulation modeling, 1477
African fruit fly,
  Ceratitis cosyra, 2068
  fecal matter, 2068
  host discrimination, 2068
  host-marking pheromone, 2068
age-stage,
  Aulacaspis yasumatsui, 1581
  Cycas revoluta, 1581
  Cycas taitungensis, 1581
  two-sex life table, 1581
aggregation,
  lesser mealworm, 2196
  pheromone, 2196
  pitfall, 2196
  poultry, 2196
aggregation pheromone, 923, 2029
  Ips, 923
  reproductive isolation, 923
  Scolytidae, 923
  semiochemical, 2029
  2-(undecyloxy)-ethanol, 2029
Agrilus planipennis, 429, 792, 1929
  ash tree, 1929
  autodissemination, 1929
  dark green trap, 429
  entomopathogenic fungus, 1929
  Eulophidae, 792
  fitness, 792
  invasive insect, 1929
  manuka oil, 429
  parasitoid, 792
  phoebe oil, 429
  Tetrastichus planipennisi, 792
  (3Z)-hexenol, 429
alate,
  drywood termite, 2213
  light wavelength, 2213
  phototropic, 2213
alfalfa,
  fluctuating thermal regime, 14
  pollinator, 14
  spring incubation, 14
alfalfa weevil,
  degree-day, 102
  integrated pest management (IPM), 102
  sheep, 102
  targeted grazing, 102
Allee effect,
  biological invasion, 1
  density dependence, 1
  eradication, 1
  invasive species, 1
Allium cepa,
  leaf color, 632
  olfactometer, 632
  onion resistance, 632
  Thrips tabaci, 632
Alphitobius diaperinus,
  cyfluthrin, 979
  diatomaceous earth, 979
  field efficacy, 979
  spinosad, 979
aluminum phosphide,
  fumigant, 135
  Lyctus africanus, 135
  Mangifera indica, 135
  Pinus roxburghii, 135
aluminum sheet bin,
  Callosobruchus chinensis, 1095
  Cicer arietinum, 1095
  jute bag, 1095
  natural product, 1095

Amblyomma,
biological control, 1979
Ixodiphagus, 1979
Rhipicephalus sanguineus, 1979
tick, 1979
ambulatory behavior,
behavioral resistance, 2187
fumigation, 2187
ozone susceptibility, 2187
stored product protection, 2187
amitraz,
collar, 2207
neckband, 2207
tick control, 2207
white-tailed deer, 2207
ammonium acetate,
Ceratitis capitata, 471
fruit fly management, 471
spinosad, 471
trimethylamine, 471
Amyelois transitella,
cytochrome P450 monooxygenase, 410
insecticide, 410
myristicin, 410
synergist, 410
Anastrepha,
areawide pest management, 823
climate, 823
population dynamics, 823
time series analysis, 823
Anastrepha obliqua,
COI, 2147
mitochondrial DNA, 2147
sequencing, 2147
West Indian fruit fly, 2147
Anoplophora glabripennis,
Acer platanoides, 2015
fecundity, 2015
longevity, 2015
neonicotinoid, 2015
antagonism,
cumulative exposure, 85
ecotoxicology, 85
mixture toxicity, 85
pesticide, 85
antibiosis, 252, 2167, 2180
antixenosis, 2180
aphid resistance, 2167
host plant resistance, 252
integrated pest management (IPM), 252
P746, 2167
single dominant gene, 2167
soybean, 2167
tolerance, 2180
wheat, 2180
wheat curl mite, 2180
antibiotic resistance,
canola cultivar, 2172
demographic parameter, 2172
Helicoverpa armigera, 2172
life table, 2172
antixenosis, 1465, 2180
antibiosis, 2180
electrical penetration graph, 1465
feeding behavior, 1465
plant resistance, 1465
tolerance, 2180
wheat, 2180
wheat curl mite, 2180
Aphelinus mali,
color, 1342
orientation, 1342
trap, 1342
aphid, 783, 1844, 1909
DNA barcoding, 1909
interference, 783
no-choice, 1844
performance, 1844
potato virus Y, 1909
Potyvirus-transmission, 783
probing, 1844
resistance, 1844
specificity, 783
vector, 1909
aphid behavior,
acquisition, 1903
artificial diet, 1903
potato virus Y, 1903
aphid management strategy,
Aphis glycines, 608
genotype, 608
Glycine max, 608
host plant resistance, 608
aphid resistance,
antibiosis, 2167
P746, 2167
single dominant gene, 2167
soybean, 2167
Aphis glycines, 25, 608, 1432, 1835
aphid management strategy, 608
Diabrotica virgifera virgifera, 25
gene pyramid, 1835
genotype, 608
Glycine max, 608
host plant resistance, 25, 608, 1835
housekeeping gene, 1432
normalization, 1432
quantitative real-time polymerase chain reaction, 1432
Rag genes, 25
rotation resistance, 25
soybean, 1835
Aphis spiraecola,
genetic differentiation, 583
genetic diversity, 583
microsatellite, 583
apical-meristem injury,
cranberry, 1366
Dasineura oxycoccana, 1366
Ericaceae, 1366
tipworm, 1366
Apion spp.,
biological control, 1620
Spintherus dubius, 1620
thiacloprid, 1620
Trifolium pratense, 1620
Apis mellifera, 313, 777, 1145, 1487, 1878, 1895
AFB, 1487
Ascosphaera apis, 1878
chalkbrood, 1878
comb orientation, 777
crop pollination, 313
fatty acid, 1145
flax oil, 1145
genetic resistance, 313
honey bee, 1145, 1487, 1878
honey harvesting, 777
lysozyme, 1878
matrix, 1487
migratory beekeeping, 313
miticide, 1895
nestmate recognition, 1145
nest pattern, 777
pesticide, 1895
polymerase chain reaction (PCR), 1487
queen, 1895
traditional hive, 777
Varroa destructor, 313, 1895
Apis mellifera anatoliaca,
acute toxicity, 1890
flumethrin, 1890
oral, 1890
Apolygus lucorum, 1603, 1921
artificial diet, 1921
early season host plant, 1603
mirid, 1921
pest management strategy, 1603
rearing, 1921
successive generations, 1921
apple, 297, 1540, 1712
Carposina sasakii, 1540
CATTS, 1540
codling moth, 297
Cydia pomonella, 297
diapause, 297
Grapholita molesta, 1712
heat shock, 1540
integrated pest management (IPM), 1712
oriental fruit moth, 1712
peach, 1712

quarantine, 297
application,
grasshopper, 1915
IGR, 1915
microsporidia, 1915
areawide,
ecology, 1597
Eoreuma, 1597
host plant, 1597
Saccharum, 1597
areawide pest management, 823, 1268
Anastrepha, 823
Bactericera cockerelli, 1268
climate, 823
insect vector, 1268
Liberibacter, 1268
population dynamics, 823
time series analysis, 823
zebra chip, 1268
Armadillidium vulgare,
Glycine max, 1629
isopod, 1629
residue management, 1629
soybean, 1629
armored scale,
avocado, 497
detection, 497
invasive species, 497
surveillance, 497
artificial diet, 802, 1903, 1921
acquisition, 1903
aphid behavior, 1903
Apolygus lucorum, 1921
biological control, 802
Coleoptera, 802
mass rearing, 802
mirid, 1921
parasitoid wasp, 802
potato virus Y, 1903
rearing, 1921
successive generations, 1921
Ascosphaera apis,
Apis mellifera, 1878
chalkbrood, 1878
honey bee, 1878
lysozyme, 1878
ash tree,
Agrilus planipennis, 1929
autodissemination, 1929
entomopathogenic fungus, 1929
invasive insect, 1929
Asian longhorned beetle,
bark beetle, 1645
emerald ash borer, 1645
exotic insect, 1645
wood borer, 1645
Asian needle ant,
control, 1751
granular bait, 1751
hydramethylnon, 1751
Pachycondyla chinensis, 1751
Atherigona soccata,
biochemical mechanisms of resistance, 1069
flavonoids, 1069
induced resistance, 1069
Sorghum bicolor, 1069
attack phase,
mass attack, 140
mass trapping, 140
semiochemical release rate, 140
synthetic lure, 140
attractant,
integrated pest management (IPM), 1335
kairomone, 1335
monitoring, 1335
oviposition, 1335
attracticidal sphere,
behavioral control, 1548
integrated pest management (IPM), 1548
Rhagoletis, 1548
spinosad, 1548
attraction,
emergence, 461
exotic species, 461
laurel wilt, 461
Scolytinae, 461
Aulacaspis yasumatsui,
age-stage, 1581
Cycas revoluta, 1581
Cycas taitungensis, 1581
two-sex life table, 1581
autodissemination,
Agrilus planipennis, 1929
ash tree, 1929
entomopathogenic fungus, 1929
invasive insect, 1929
avermectin,
acetamiprid, 988
Brontispa longissimi, 988
β-cypermethrin, 988
insecticide susceptibility, 988
avocado, 402, 497, 659, 1178, 2035
armored scale, 497
α-copaene, 659
detection, 497
developmental biology, 2035
Diaspididae, 2035
host plant resistance, 1178
invasive species, 497, 2035
laurel wilt, 659
life table, 402
Mediterranean fruit fly, 1178
Perú, 402
quarantine security, 1178
Redbay ambrosia beetle, 659
risk analysis, 1178
sesquiterpene, 659
sex pheromone, 402
Stenoma catenifer, 402
surveillance, 497

B

Bacillus thuringiensis (*Bt*), 214, 616, 732, 767, 1164, 1773, 1863, 2122
adult mortality, 1863
baseline susceptibility, 214
camptothecin (CPT), 1164
Cnaphalocrocis medinalis, 2122
corn borer, 214
Cry toxin, 2122
Diabrotica, 767
diet, 1863
genetically engineered crop, 767
Helicoverpa zea, 1773
Hemiptera, 616
insect histopathology, 1164
larval mortality, 732
Lygus, 616
nucleopolyhedrovirus (NPV), 1164
Ostrinia nubilalis, 1773
refuge, 767, 1773
resistance, 214, 767
selectivity, 2122
Stomoxys calcitrans, 732, 1863
susceptibility, 2122
synergism, 1164
tarnished plant bug, 616
temperature, 732
transgenic cotton, 616
transgenic maize, 214
Bactericera cockerelli,
areawide pest management, 1268
insect vector, 1268
Liberibacter, 1268
zebra chip, 1268
Bactrocera invadens,
cold disinfestation, 1963
commodity treatment, 1963
phytosanitary, 1963
quarantine, 1963
Bactrocera oleae,
adult survival, 186
humidity, 186
reproductive biology, 186
temperature, 186
Bactrocera tryoni,
domestication, 1051
genetic drift, 1051
mass rearing, 1051
selection, 1051
baculovirus, 344, 1503
geographic virus isolate, 344
Lymantria dispar asiatica, 344
M. bicoloratus, 1503
parasitoid, 1503

pathogenicity, 344
S. exigua, 1503
S. exigua multiple nucleopolyhedrovirus, 1503
virulence, 344
bait,
boric acid, 1385
silverfish, 1385
toxicity, 1385
baiting system,
cardboard, 1758
Coptotermes formosanus, 1758
maize cob, 1758
pine wood, 1758
bark beetle, 1640, 1645
Asian longhorned beetle, 1645
blue-stained sapwood, 1640
emerald ash borer, 1645
exotic insect, 1645
Formosan subterranean termite, 1640
ophiostomatoid fungi, 1640
southern yellow pine, 1640
wood borer, 1645
Bartlett's sequential test,
binomial sampling, 1077
generalized linear mixed model, 1077
baseline data,
basic model, 1024
50% lethal dose, 1024
proportionality, 1024
weight study, 1024
baseline susceptibility,
Bacillus thuringiensis (*Bt*), 214
orn borer, 214
resistance, 214
transgenic maize, 214
basic model,
baseline data, 1024
50% lethal dose, 1024
proportionality, 1024
weight study, 1024
Bayesian statistics,
Abbott's formula, 1856
EC_{50}, 1856
range estimation, 1856
repellency, 1856
Beauveria bassiana,
botanical, 2097
IGR, 2097
insecticide, 2097
survival, 2097
bed bug, 1789, 2107
Cimex lectularius, 1789, 2107
desiccant dust, 1789
management, 2107
pyrethroid, 1789
repellency, 2107
resistance, 1789
ultrasound, 2107
Begomovirus,
host plant resistance, 1447
transmission, 1447
vector, 1447
behavioral control,
attracticidal sphere, 1548
integrated pest management (IPM), 1548
Rhagoletis, 1548
spinosad, 1548
behavioral resistance,
ambulatory behavior, 2187
fumigation, 2187
ozone susceptibility, 2187
stored product protection, 2187
Bemisia argentifolii,
microsatellite marker, 753
mitochondrial cytochrome oxidase I subunit, 753
New World biotype, 753
Bemisia tabaci,
diagnostic technique, 242
genetic group, 242
mitochondrial cytochrome oxidase I gene, 242
polymerase chain reaction-restriction fragment length polymorphism, 242
Bemisia tabaci biotype B,
biological invasion, 930
biology, 930
biotype Q, 930
host plant, 930
binomial sampling,
Bartlett's sequential test, 1077
generalized linear mixed model, 1077
bioassay,
Grapholita molesta, 1426
insecticide resistance, 1426
joint action, 1426
biochemical mechanisms of resistance,
Atherigona soccata, 1069
flavonoids, 1069
induced resistance, 1069
Sorghum bicolor, 1069
bioclimatic modeling, 739, 1115
cost of management, 1115
diamondback moth, 1115
integrated pest management (IPM), 739, 1115
pest control, 1115
Plutella xylostella, 739
population outbreak, 739
resistance, 739
biodegradation,
escape behavior, 62
hypoxia, 62
pig manure, 62
separation, 62
biodemographic study,
Ceratitis rosa, 371
fruit fly, 371
longevity, 371
Tephritidae, 371
biofuel feedstock,
fertilization, 878
herbivory, 878
Schistocerca americana, 878
Spodoptera frugiperda, 878
biological control, 92, 161, 354, 386, 802, 1311, 1509, 1520, 1532, 1620, 1979
Amblyomma, 1979
Apion spp., 1620
artificial diet, 802
chemical control, 92
Coccinellidae, 1509
Coleoptera, 802
diamondback moth, 354
ectoparasitoid, 161
egg parasitoid, 92, 1532
egg structure, 1520
exotic invasive pest, 1311
glassy-winged sharpshooter, 1532
Hemiptera, 1509
host genotype, 354
identification, 1520
insecticide, 354
interference competition, 386
Ixodiphagus, 1979
leafminer, 161
mass rearing, 802
native natural enemy, 1311
Necremnus spp., 1311
operculum, 1520
Orius, 1520
parasitoid biology, 1532
parasitoid wasp, 802
predator, 1509
Pseudacteon tricuspis, 386
resistance, 354
Rhipicephalus sanguineus, 1979
sex ratio, 161, 386
social form, 386
Spintherus dubius, 1620
thiacloprid, 1620
tick, 1979
Trichogramma japonicum, 92
Trifolium pratense, 1620
tritrophic interactions, 161
biological invasion, 1, 438, 930
Allee effect, 1
Bemisia tabaci biotype B, 930
biology, 930
biotype Q, 930
density dependence, 1
eradication, 1
firewood, 438
host plant, 930

human-mediated dispersal, 438
invasive forest pest, 438
invasive species, 1
long-distance dispersal, 438
biological parameter,
fecundity, 1809
host plant resistance, 1809
potato cultivar, 1809
potato tuberworm, 1809
biology,
Bemisia tabaci biotype B, 930
biological invasion, 930
biotype Q, 930
host plant, 930
biophysical model,
degree-day model, 860
temperature-dependent development, 860
western tarnished plant bug, 860
biorational,
meliponini, 1234
pesticide, 1234
Saccharopolyspora spinosa, 1234
social, 1234
biotype, 232, 1816
DNA methylation, 232
host-plant resistance, 1816
salivary gland, 232
sodium bisulfite, 232
thrip *Frankliniella occidentalis*, 1816
virulence, 232
biotype Q,
Bemisia tabaci biotype B, 930
biological invasion, 930
biology, 930
host plant, 930
bird cherry-oat aphid,
plant resistance categories, 592
wheat line, 592
blacklegged tick,
entomopathogenic fungi, 222
Ixodes scapularis, 222
Metarhizium anisopliae, 222
tick control, 222
black soldier fly,
fermentation, 374
Hermetia illucens, 374
leachate processing, 374
blended-seed refuge,
Bt corn, 1214
dispersal, 1214
European corn borer, 1214
resistance management, 1214
blow fly,
Calliphora nigribarbis, 1796
highly pathogenic avian influenza, 1796
insecticide-treated netting screen, 1796
permethrin, 1796
blue-stained sapwood,
bark beetle, 1640
Formosan subterranean termite, 1640
ophiostomatoid fungi, 1640
southern yellow pine, 1640
Bombus,
honey bee, 1130
Megachile, 1130
non-*Apis* bee, 1130
Osmia, 1130
Bombus ephippiatus,
bumble bee, 1871
greenhouse, 1871
pollination, 1871
tomato, 1871
boric acid, 1139, 1385
bait, 1385
fipronil, 1139
indoxacarb, 1139
silverfish, 1385
survival, 1139
toxic bait efficacy, 1139
toxicity, 1385
botanical,
Beauveria bassiana, 2097
IGR, 2097
insecticide, 2097
survival, 2097
botanical insecticide, 1329, 2115
Camptomyia corticalis, 1329
cecidomyiid gall midge, 1329
chloroform extract, 2115
essential oil, 1329
natural fumigant, 1329
subterranean termite, 2115
Brontispa longissimi,
acetamiprid, 988
avermectin, 988
β-cypermethrin, 988
insecticide susceptibility, 988
brood pheromone, 304, 1134
colony growth, 1134
honey, 304
honey bee, 304, 1134
nutrition, 1134
protein supplement, 1134
SuperBoost, 304
survival, 304
Bt corn,
blended-seed refuge, 1214
dispersal, 1214
European corn borer, 1214
resistance management, 1214
Bt maize,
Diabrotica virgifera virgifera, 1248
maize, 1248
nontarget, 1248
pyramid, 1248
bug bomb,
Cimex lectularius, 957
fogger, 957
pyrethroid, 957
resistance, 957
bumble bee, 34, 1871
Bombus ephippiatus, 1871
foraging tradeoff, 34
greenhouse, 34, 1871
pollination, 34, 1871
tomato, 34, 1871

C

C. quinquefasciatus,
esterases resistance, 971
insecticide, 971
naled, 971
resmethrin, 971
cage,
Reunion Island, 1358
spinosad, 1358
Synéis-appât, 1358
Tephritidae, 1358
Calliphora nigribarbis,
blow fly, 1796
highly pathogenic avian influenza, 1796
insecticide-treated netting screen, 1796
permethrin, 1796
Callosobruchus chinensis,
aluminum sheet bin, 1095
Cicer arietinum, 1095
jute bag, 1095
natural product, 1095
Camptomyia corticalis,
botanical insecticide, 1329
cecidomyiid gall midge, 1329
essential oil, 1329
natural fumigant, 1329
camptothecin (CPT),
Bacillus thuringiensis (*Bt*), 1164
insect histopathology, 1164
nucleopolyhedrovirus (NPV), 1164
synergism, 1164
Candidatus Phytoplasma mali,
immigration dynamic, 322
overwintered adult, 322
psyllid vector, 322
temperature, 322
canola,
crop rotation, 1261
Delia spp., 1261
root maggot, 1261
canola cultivar,
antibiotic resistance, 2172
demographic parameter, 2172
Helicoverpa armigera, 2172
life table, 2172

carbamate,
acetylcholinesterase insensitivity, 540
general esterase, 540
glutathione-*S*-transferase, 540
huanglongbing, 540
carboxylesterase,
cytochrome P450, 1034
glutathione transferase, 1034
quercetin, 1034
silkworm, 1034
cardboard,
baiting system, 1758
Coptotermes formosanus, 1758
maize cob, 1758
pine wood, 1758
Cardinium,
CI, 2161
fitness, 2161
sex ratio, 2161
Sogatella furcifera, 2161
Carposina sasakii,
apple, 1540
CATTS, 1540
heat shock, 1540
CATTS,
apple, 1540
Carposina sasakii, 1540
heat shock, 1540
cecidomyiid gall midge,
botanical insecticide, 1329
Camptomyia corticalis, 1329
essential oil, 1329
natural fumigant, 1329
Cerambycidae, 395, 837, 906
flight-intercept trap, 837
Fluon, 395
fuscumol acetate, 906
2,3-hexanediol, 906
3-hydroxy-2-hexanone, 906
inhibition, 906
monitoring, 837
pheromone trap, 395
trapping, 837
wood-borer detection, 837
wood-boring insect, 395
Ceratitis capitata,
ammonium acetate, 471
fruit fly management, 471
spinosad, 471
trimethylamine, 471
Ceratitis cosyra,
African fruit fly, 2068
fecal matter, 2068
host discrimination, 2068
host-marking pheromone, 2068
Ceratitis rosa,
biodemographic study, 371
fruit fly, 371
longevity, 371
Tephritidae, 371
chalkbrood,
Apis mellifera, 1878
Ascosphaera apis, 1878
honey bee, 1878
lysozyme, 1878
chemical control, 92, 1998
biological control, 92
egg parasitoid, 92
Eoreuma loftini, 1998
neonate, 1998
novaluron, 1998
sugarcane, 1998
Trichogramma japonicum, 92
cherelle wilt,
cocoa mirids, 1285
cocoa yield, 1285
Sahlbergella singularis, 1285
Theobroma cacao, 1285
chlorantraniliprole, 549, 866, 1019, 1736
eastern subterranean termite, 1736
flubendiamide, 866
horizontal transfer, 1736
insecticide, 1736
Leptinotarsa decemlineata, 549
novaluron, 866
Plutella xylostella, 1019
resistance monitoring, 1019
Reticulitermes flavipes, 1736
spinetoram, 866
spirotetramat, 866
synergist, 549
toxicity, 549
chloroform extract,
botanical insecticide, 2115
subterranean termite, 2115
chrysopid,
cypermethrin, 1982
endosulfan, 1982
methoxyfenozide, 1982
spinosad, 1982
CI,
Cardinium, 2161
fitness, 2161
sex ratio, 2161
Sogatella furcifera, 2161
Cicadulina bipunctata
Forage grass, 129
Laodelphax striatellus, 129
N-formylloline, 129
Sogatella furcifera, 129
Cicer arietinum,
aluminum sheet bin, 1095
Callosobruchus chinensis, 1095
jute bag, 1095
natural product, 1095
Cimex lectularius, 957, 1789, 2107
bed bug, 1789, 2107
bug bomb, 957
desiccant dust, 1789
fogger, 957
management, 2107
pyrethroid, 957, 1789
repellency, 2107
resistance, 957, 1789
ultrasound, 2107
citronellal,
eugenol, 72
flight muscles, 72
pulegone, 72
α-terpineol, 72
citrus,
life cycle, 338
net reproductive rate, 338
parasitism, 338
Citrus limon,
Mexican fruit fly, 363
Rutaceae, 363
Tephritidae, 363
citrus orchard,
GF-120, 1277
hydrolyzed-protein bait, 1277
monitoring, 1277
phenology, 1277
climate,
Anastrepha, 823
areawide pest management, 823
population dynamics, 823
time series analysis, 823
Cnaphalocrocis medinalis,
Bacillus thuringiensis (*Bt*), 2122
Cry toxin, 2122
selectivity, 2122
susceptibility, 2122
coating,
essential oil, 1238
Frankliniella occidentalis, 1238
polymer, 1238
thrip, 1238
Coccinellidae,
biological control, 1509
Hemiptera, 1509
predator, 1509
cocoa mirids,
cherelle wilt, 1285
cocoa yield, 1285
Sahlbergella singularis, 1285
Theobroma cacao, 1285
cocoa yield,
cherelle wilt, 1285
cocoa mirids, 1285
Sahlbergella singularis, 1285
Theobroma cacao, 1285
Cocos nucifera,
method, 1419
pesticide, 1419
standardization, 1419
codling moth, 297, 872
apple, 297
Cydia pomonella, 297
diapause, 297

fitness trade-off, 872
insecticide resistance, 872
quarantine, 297
COI,
Anastrepha obliqua, 2147
mitochondrial DNA, 2147
sequencing, 2147
West Indian fruit fly, 2147
cold disinfestation,
Bactrocera invadens, 1963
commodity treatment, 1963
phytosanitary, 1963
quarantine, 1963
Coleoptera, 802, 2076
artificial diet, 802
biological control, 802
isopropanol, 2076
kairomone, 2076
mass rearing, 802
parasitoid wasp, 802
Scarabaeidae, 2076
collar,
amitraz, 2207
neckband, 2207
tick control, 2207
white-tailed deer, 2207
colony elimination, 523, 533
colony suppression, 523
microsatellite, 523
noviflumuron bait, 533
Reticulitermes flavipes, 523
Reticulitermes hageni, 523
subterranean termite, 533
termite baiting, 533
colony growth,
brood pheromone, 1134
honey bee, 1134
nutrition, 1134
protein supplement, 1134
colony suppression,
colony elimination, 523
microsatellite, 523
Reticulitermes flavipes, 523
Reticulitermes hageni, 523
color,
Aphelinus mali, 1342
orientation, 1342
trap, 1342
comb orientation,
Apis mellifera, 777
honey harvesting, 777
nest pattern, 777
traditional hive, 777
commodity treatment,
Bactrocera invadens, 1963
cold disinfestation, 1963
phytosanitary, 1963
quarantine, 1963
common smut,
corn earworm, 1457
maize weevil, 1457
principal components analysis, 1457
stink bug–discolored kernel, 1457
community index,
intercropping, 854
natural enemy, 854
Sitobion avenae, 854
wheat, 854
concrete,
methoprene, 718
residual efficacy, 718
surface treatment, 718
varnished wood, 718
cone insect,
Dioryctria abietella, 916
Eupithecia abietaria, 916
Norway spruce, 916
seed insect, 916
conservation biological control,
natural enemy, 490
potato psyllid, 490
tomato psyllid, 490
toxicity, 490
control,
Asian needle ant, 1751
granular bait, 1751
hydramethylnon, 1751
Pachycondyla chinensis, 1751
controlled atmosphere,
disinfestation, 1171
ethanol, 1171
methyl bromide alternative, 1171
modified atmosphere, 1171
α-copaene,
avocado, 659
laurel wilt, 659
Redbay ambrosia beetle, 659
sesquiterpene, 659
copper,
development, 288
population dynamics, 288
Spodoptera litura, 288
Coptotermes formosanus, 1758, 1766
baiting system, 1758
cardboard, 1758
dust formulation, 1766
maize cob, 1758
pine wood, 1758
termite control, 1766
termiticide, 1766
corn borer,
Bacillus thuringiensis (*Bt*), 214
baseline susceptibility, 214
resistance, 214
transgenic maize, 214
corn earworm,
common smut, 1457
maize weevil, 1457
principal components analysis, 1457
stink bug–discolored kernel, 1457
cost of management,
bioclimatic modeling, 1115
diamondback moth, 1115
integrated pest management (IPM), 1115
pest control, 1115
cotton, 2007, 2136
Lygus, 2007
microsatellite, 2136
pest management, 2136
resistance, 2136
sampling, 2007
soybean, 2136
tarnished plant bug, 2007
threshold, 2007
cotton fleahopper,
Creontiades signatus, 896
pest sampling, 896
Pseudatomoscelis seriatus, 896
verde plant bug, 896
cotton with colored fibers,
damage, 113
fennel, 113
Hyadaphis foeniculi, 113
intercropping, 113
cranberry,
apical-meristem injury, 1366
Dasineura oxycoccana, 1366
Ericaceae, 1366
tipworm, 1366
Creontiades signatus,
cotton fleahopper, 896
pest sampling, 896
Pseudatomoscelis seriatus, 896
verde plant bug, 896
crop pollination,
Apis mellifera, 313
genetic resistance, 313
migratory beekeeping, 313
Varroa destructor, 313
crop rotation,
canola, 1261
Delia spp., 1261
root maggot, 1261
crucifer,
field predation, 1719
grower recommendation, 1719
spined soldier bug, 1719
yellowmargined leaf beetle, 1719
Cry toxin,
Bacillus thuringiensis (*Bt*), 2122
Cnaphalocrocis medinalis, 2122
selectivity, 2122
susceptibility, 2122

cryptic species,
Dasineura oxycoccana, 1205
phenology, 1205
Vaccinium corymbosum, 1205
Vaccinium macrocarpon, 1205
cultivar,
Nesidiocoris tenuis, 2085
release strategy, 2085
Trichogramma achaeae, 2085
Tuta absoluta, 2085
cumulative exposure,
antagonism, 85
ecotoxicology, 85
mixture toxicity, 85
pesticide, 85
Cycas revoluta,
age-stage, 1581
Aulacaspis yasumatsui, 1581
Cycas taitungensis, 1581
two-sex life table, 1581
Cycas taitungensis,
age-stage, 1581
Aulacaspis yasumatsui, 1581
Cycas revoluta, 1581
two-sex life table, 1581
Cyclocephala,
Paenibacillus, 1320
Popillia japonica, 1320
soil insect, 1320
Tiphiidae, 1320
Cydia pomonella,
apple, 297
codling moth, 297
diapause, 297
quarantine, 297
cyfluthrin,
Alphitobius diaperinus, 979
diatomaceous earth, 979
field efficacy, 979
spinosad, 979
Cylas formicarius,
Diabrotica, 651
Ipomoea, 651
plant resistance, 651
white grub, 651
cypermethrin,
chrysopid, 1982
endosulfan, 1982
methoxyfenozide, 1982
spinosad, 1982
β-cypermethrin,
acetamiprid, 988
avermectin, 988
Brontispa longissimi, 988
insecticide susceptibility, 988
cytochrome oxidase I, 418, 573
fall armyworm, 573
haplotype, 418
host race, 573
PCR-restriction fragment-length polymorphism, 573
South America, 573
Spodoptera frugiperda, 418
triose phosphate isomerase, 418
cytochrome P450, 1006, 1034
acrinathrin, 1006
carboxylesterase, 1034
glutathione transferase, 1034
insecticide resistance, 1006
over-expression, 1006
quercetin, 1034
silkworm, 1034
western flower thrip, 1006
cytochrome P450 monooxygenase, 410, 560
Amyelois transitella, 410
insecticide, 410
insecticide induction, 560
myristicin, 410
psocid, 560
quantitative polymerase chain reaction, 560
synergist, 410

D

damage,
cotton with colored fibers, 113
fennel, 113
Hyadaphis foeniculi, 113
intercropping, 113
dark green trap,
Agrilus planipennis, 429
manuka oil, 429
phoebe oil, 429
(3Z)-hexenol, 429
Dasineura oxycoccana, 1205, 1366
apical-meristem injury, 1366
cranberry, 1366
cryptic species, 1205
Ericaceae, 1366
phenology, 1205
tipworm, 1366
Vaccinium corymbosum, 1205
Vaccinium macrocarpon, 1205
date palm pest,
light trap, 2062
Oryctes agamemnon arabicus, 2062
photoperiod, 2062
rhinoceros beetle, 2062
degree-day, 102, 1573
alfalfa weevil, 102
developmental rate, 1573
Herpetogramma phaeopteralis, 1573
integrated pest management (IPM), 102
nonlinear model, 1573
sheep, 102
targeted grazing, 102
turfgrass, 1573
degree-day model,
biophysical model, 860
temperature-dependent development, 860
western tarnished plant bug, 860
Delia spp.,
canola, 1261
crop rotation, 1261
root maggot, 1261
delimitation,
detection, 272
emerald ash borer, 272
invasive species, 272
surveillance, 272
demographic parameter,
antibiotic resistance, 2172
canola cultivar, 2172
Helicoverpa armigera, 2172
life table, 2172
Dendroctonus ponderosae,
acetophenone, 149
nonhost angiosperm volatile, 149
pest management, 149
Pinus ponderosa, 149
density dependence,
Allee effect, 1
biological invasion, 1
eradication, 1
invasive species, 1
desiccant dust,
bed bug, 1789
Cimex lectularius, 1789
pyrethroid, 1789
resistance, 1789
detection, 272, 497, 1350, 1557
armored scale, 497
avocado, 497
delimitation, 272
Drosophila suzukii, 1350
emerald ash borer, 272
eradication program, 1557
fruit fly, 1557
invasive species, 272, 497
monitoring, 1350
surveillance, 272, 497
survey, 1557
trapping, 1350
deterrent,
diamondback moth, 1379
lemon-scented tea tree oil (LSO), 1379
repellant, 1379
development, 288, 964, 1471
copper, 288
eclosion, 964
hatching, 964
mating, 964
nutritional quality, 1471
population dynamics, 288
reproduction, 1471

Spodoptera litura, 288
stored product, 1471
Tribolium castaneum, 1471
vegetative insecticidal protein, 964
developmental biology,
avocado, 2035
Diaspididae, 2035
invasive species, 2035
developmental rate,
degree-day, 1573
Herpetogramma phaeopteralis, 1573
nonlinear model, 1573
turfgrass, 1573
Diabrotica, 651, 767
Bacillus thuringiensis (*Bt*), 767
Cylas formicarius, 651
genetically engineered crop, 767
Ipomoea, 651
plant resistance, 651
refuge, 767
resistance, 767
white grub, 651
Diabrotica virgifera virgifera, 25, 1248, 1407
Aphis glycines, 25
Bt maize, 1248
feeding behavior, 1407
host plant resistance, 25
maize, 1248
nontarget, 1248
pest resistance, 1407
pyramid, 1248
Rag genes, 25
rotation resistance, 25
transgenic maize, 1407
diagnostic technique,
Bemisia tabaci, 242
genetic group, 242
mitochondrial cytochrome oxidase I gene, 242
polymerase chain reaction-restriction fragment length polymorphism, 242
diamondback moth, 354, 1115, 1379
bioclimatic modeling, 1115
biological control, 354
cost of management, 1115
deterrent, 1379
host genotype, 354
insecticide, 354
integrated pest management (IPM), 1115
lemon-scented tea tree oil (LSO), 1379
pest control, 1115
repellant, 1379
resistance, 354
diapause,
apple, 297
codling moth, 297
Cydia pomonella, 297
quarantine, 297
diapausing larvae,
low pressure, 679
methyl bromide alternative, 679
monoterpenoid, 679
Plodia interpunctella, 679
Diaphorina citri,
electrical penetration graph, 1492
feeding disruption, 1492
insect behavior, 1492
insecticide, 1492
Diaspididae,
avocado, 2035
developmental biology, 2035
invasive species, 2035
diatomaceous earth, 670, 979
Alphitobius diaperinus, 979
cyfluthrin, 979
field efficacy, 979
insecticidal potential, 670
natural zeolite, 670
Sitophilus oryzae, 670
spinosad, 979
Tribolium castaneum, 670
Diatraea grandiosella,
Diatraea saccharalis, 1825
Spodoptera exigua, 1825
Spodoptera frugiperda, 1825
transgenic corn, 1825
Diatraea saccharalis,
Diatraea grandiosella, 1825
Spodoptera exigua, 1825
Spodoptera frugiperda, 1825
transgenic corn, 1825
diet, 26, 1863
adult mortality, 1863
Aethina tumida, 26
Bacillus thuringiensis (*Bt*), 1863
population growth, 26
resource acquisition, 26
Stomoxys calcitrans, 1863
temperature, 26
digestive enzyme,
feeding performance, 1439
Helicoverpa armigera, 1439
host plant, 1439
Dioryctria abietella,
cone insect, 916
Eupithecia abietaria, 916
Norway spruce, 916
seed insect, 916
discriminating dose,
imidacloprid, 1392
immature, 1392
Planococcus ficus, 1392
resistance monitoring, 1392
disinfestation,
controlled atmosphere, 1171
ethanol, 1171
methyl bromide alternative, 1171
modified atmosphere, 1171
dispersal, 259, 1214
blended-seed refuge, 1214
Bt corn, 1214
European corn borer, 1214
forecasting, 259
migration, 259
resistance management, 1214
disruption,
Epiphyas postvittana, 1694
E11-tetradecenyl acetate, 1694
light brown apple moth, 1694
pheromone, 1694
Diuraphis noxia,
adaptation, 1057
genetic diversity, 1057
plant resistance, 1057
wheat, 1057
DNA barcoding,
aphid, 1909
potato virus Y, 1909
vector, 1909
DNA methylation,
biotype, 232
salivary gland, 232
sodium bisulfite, 232
virulence, 232
domestication,
Bactrocera tryoni, 1051
genetic drift, 1051
mass rearing, 1051
selection, 1051
Douglas-fir beetle,
MCH, 451
3-methylcylcohex-2-en-1-one, 451
pheromone plume, 451
Scolytinae, 451
Drosophila suzukii,
detection, 1350
monitoring, 1350
trapping, 1350
drywood termite,
alate, 2213
light wavelength, 2213
phototropic, 2213
dust formulation,
Coptotermes formosanus, 1766
termite control, 1766
termiticide, 1766

E

early season host plant,
Apolygus lucorum, 1603
pest management strategy, 1603

eastern subterranean termite,
chlorantraniliprole, 1736
horizontal transfer, 1736
insecticide, 1736
Reticulitermes flavipes, 1736
EC_{50},
Abbott's formula, 1856
Bayesian statistics, 1856
range estimation, 1856
repellency, 1856
(E)-β-caryophyllene,
(E)-fall armyworm, 120
maize, 120
resistance, 120
volatile, 120
eclosion,
development, 964
hatching, 964
mating, 964
vegetative insecticidal protein, 964
ecology,
areawide, 1597
Eoreuma, 1597
host plant, 1597
Saccharum, 1597
economic analysis, 196, 1612
emerald ash borer, 196
preemptive removal, 196
pyrethroid, 1612
regional survey, 1612
risk aversion, 1612
urban forest management, 196
economic injury level,
flower thrips, 480
Frankliniella bispinosa, 480
southern highbush blueberry, 480
Vaccinium corymbosum × *Vaccinium darrowi*, 480
economics,
heat treatment, 709
insect mortality, 709
sanitation, 709
temperature, 709
ecotoxicology,
antagonism, 85
cumulative exposure, 85
mixture toxicity, 85
pesticide, 85
ectoparasitoid,
biological control, 161
leafminer, 161
sex ratio, 161
tritrophic interactions, 161
efficacy assessment,
heat treatment, 703
red flour beetle, 703
sanitation, 703
temperature, 703
egg parasitoid, 92, 1532
biological control, 92, 1532
chemical control, 92
glassy-winged sharpshooter, 1532
parasitoid biology, 1532
Trichogramma japonicum, 92
egg structure,
biological control, 1520
identification, 1520
operculum, 1520
Orius, 1520
electrical penetration graph, 1465, 1492
antixenosis, 1465
Diaphorina citri, 1492
feeding behavior, 1465
feeding disruption, 1492
insect behavior, 1492
insecticide, 1492
plant resistance, 1465
electroantennogram, 890, 1149
green leaf volatile, 1149
pan trap, 890
plant volatile, 890
Plutella xylostella, 1149
sex pheromone, 890, 1149
soybean aphid, 890
trap catch, 1149
ELISA,
guttation, 67
imidacloprid, 67
melon, 67
emerald ash borer, 196, 272, 1645
Asian longhorned beetle, 1645
bark beetle, 1645
delimitation, 272
detection, 272
economic analysis, 196
exotic insect, 1645
invasive species, 272
preemptive removal, 196
surveillance, 272
urban forest management, 196
wood borer, 1645
emergence, 461, 1524
adult longevity, 1524
attraction, 461
exotic species, 461
Gryon japonicum, 1524
laurel wilt, 461
Ooencyrtus nezarae, 1524
parasitism, 1524
Scolytinae, 461
Empoasca vitis,
irrigation system, 176
vineyard, 176
water stress, 176
endosulfan,
chrysopid, 1982
cypermethrin, 1982
methoxyfenozide, 1982
spinosad, 1982
entomopathogenic fungi,
blacklegged tick, 222
Ixodes scapularis, 222
Metarhizium anisopliae, 222
tick control, 222
entomopathogenic fungus,
Agrilus planipennis, 1929
ash tree, 1929
autodissemination, 1929
invasive insect, 1929
entomopathogenic nematode,
genetically modified crop, 994
meta-analysis, 994
Pectinophora gossypiella, 994
tritrophic interaction, 994
Eoreuma,
areawide, 1597
ecology, 1597
host plant, 1597
Saccharum, 1597
Eoreuma loftini,
chemical control, 1998
neonate, 1998
novaluron, 1998
sugarcane, 1998
Epiphyas postvittana,
disruption, 1694
E11-tetradecenyl acetate, 1694
light brown apple moth, 1694
pheromone, 1694
eradication,
Allee effect, 1
biological invasion, 1
density dependence, 1
invasive species, 1
eradication program,
detection, 1557
fruit fly, 1557
survey, 1557
Ericaceae,
apical-meristem injury, 1366
cranberry, 1366
Dasineura oxycoccana, 1366
tipworm, 1366
escape behavior,
biodegradation, 62
hypoxia, 62
pig manure, 62
separation, 62
essential oil, 379, 1238, 1329
botanical insecticide, 1329
Camptomyia corticalis, 1329
cecidomyiid gall midge, 1329
coating, 1238
Frankliniella occidentalis, 1238
natural fumigant, 1329
polymer, 1238
repellency, 379
sandalwood, 379
santalol, 379
Tetranychus urticae, 379

thrip, 1238
esterases resistance,
C. quinquefasciatus, 971
insecticide, 971
naled, 971
resmethrin, 971
E11-tetradecenyl acetate,
disruption, 1694
Epiphyas postvittana, 1694
light brown apple moth, 1694
pheromone, 1694
ethanol,
controlled atmosphere, 1171
disinfestation, 1171
methyl bromide alternative, 1171
modified atmosphere, 1171
eugenol,
citronellal, 72
flight muscles, 72
pulegone, 72
α-terpineol, 72
Eulophidae,
Agrilus planipennis, 792
fitness, 792
parasitoid, 792
Tetrastichus planipennisi, 792
Eupithecia abietaria,
cone insect, 916
Dioryctria abietella, 916
Norway spruce, 916
seed insect, 916
European corn borer,
blended-seed refuge, 1214
Bt corn, 1214
dispersal, 1214
resistance management, 1214
exotic insect,
Asian longhorned beetle, 1645
bark beetle, 1645
emerald ash borer, 1645
wood borer, 1645
exotic invasive pest,
biological control, 1311
native natural enemy, 1311
Necremnus spp., 1311
exotic species,
attraction, 461
emergence, 461
laurel wilt, 461
Scolytinae, 461

F

fall armyworm, 120, 573
cytochrome oxidase I, 573
(E)-β-caryophyllene, 120
host race, 573
maize, 120
PCR-restriction fragment-length polymorphism, 573
resistance, 120
South America, 573
volatile, 120
fatty acid,
Apis mellifera, 1145
flax oil, 1145
honey bee, 1145
nestmate recognition, 1145
fecal matter,
African fruit fly, 2068
Ceratitis cosyra, 2068
host discrimination, 2068
host-marking pheromone, 2068
fecundity, 1809, 2015
Acer platanoides, 2015
Anoplophora glabripennis, 2015
biological parameter, 1809
host plant resistance, 1809
longevity, 2015
neonicotinoid, 2015
potato cultivar, 1809
potato tuberworm, 1809
feeding behavior, 1407, 1465
antixenosis, 1465
Diabrotica virgifera virgifera, 1407
electrical penetration graph, 1465
pest resistance, 1407
plant resistance, 1465
transgenic maize, 1407
feeding disruption,
Diaphorina citri, 1492
electrical penetration graph, 1492
insect behavior, 1492
insecticide, 1492
feeding preference,
intraplant movement, 847
pest management, 847
sampling, 847
stink bug, 847
feeding performance,
digestive enzyme, 1439
Helicoverpa armigera, 1439
host plant, 1439
fennel,
cotton with colored fibers, 113
damage, 113
Hyadaphis foeniculi, 113
intercropping, 113
fermentation,
black soldier fly, 374
Hermetia illucens, 374
leachate processing, 374
Ferrisia,
mealybug, 1681
pistachio, 1681
Pistacia, 1681
Pseudococcidae, 1681
fertilization,
biofuel feedstock, 878
herbivory, 878
Schistocerca americana, 878
Spodoptera frugiperda, 878
field efficacy,
Alphitobius diaperinus, 979
cyfluthrin, 979
diatomaceous earth, 979
spinosad, 979
field predation,
crucifer, 1719
grower recommendation, 1719
spined soldier bug, 1719
yellowmargined leaf beetle, 1719
fipronil,
boric acid, 1139
indoxacarb, 1139
survival, 1139
toxic bait efficacy, 1139
fipronil resistance,
GABA receptor, 1781
Oulema oryzae Kuwayama, 1781
rice leaf beetle, 1781
firewood,
biological invasion, 438
human-mediated dispersal, 438
invasive forest pest, 438
long-distance dispersal, 438
fitness, 792, 2161
Agrilus planipennis, 792
Cardinium, 2161
CI, 2161
Eulophidae, 792
parasitoid, 792
sex ratio, 2161
Sogatella furcifera, 2161
Tetrastichus planipennisi, 792
fitness cost,
acetamiprid, 1401
resistance, 1401
whitefly, 1401
fitness trade-off,
codling moth, 872
insecticide resistance, 872
flavonoids,
Atherigona soccata, 1069
biochemical mechanisms of resistance, 1069
induced resistance, 1069
Sorghum bicolor, 1069
flax oil,
Apis mellifera, 1145
fatty acid, 1145
honey bee, 1145
nestmate recognition, 1145
flight-intercept trap,
Cerambycidae, 837
monitoring, 837
trapping, 837
wood-borer detection, 837

flight muscles,
citronellal, 72
eugenol, 72
pulegone, 72
α-terpineol, 72
floriculture,
greenhouse, 505
light intensity, 505
pest management, 505
systemic insecticide, 505
flour mill,
fumigation, 686
integrated pest management (IPM), 686
spatial distribution, 686
Tribolium castaneum, 686
flower thrips,
economic injury level, 480
Frankliniella bispinosa, 480
southern highbush blueberry, 480
Vaccinium corymbosum × *Vaccinium darrowi*, 480
flubendiamide,
Chlorantraniliprole, 866
novaluron, 866
spinetoram, 866
spirotetramat, 866
fluctuating thermal regime,
alfalfa, 14
pollinator, 14
spring incubation, 14
flumethrin,
acute toxicity, 1890
Apis mellifera anatoliaca, 1890
oral, 1890
Fluon,
Cerambycidae, 395
pheromone trap, 395
wood-boring insect, 395
fogger,
bug bomb, 957
Cimex lectularius, 957
pyrethroid, 957
resistance, 957
Forage grass,
Cicadulina bipunctata
Laodelphax striatellus, 129
N-formylloline, 129
Sogatella furcifera, 129
foraging tradeoff,
bumble bee, 34
greenhouse, 34
pollination, 34
tomato, 34
forecasting,
dispersal, 259
migration, 259
Formosan subterranean termite, 518, 1640, 1746
bark beetle, 1640
blue-stained sapwood, 1640
Hurricane Katrina, 518
inundation, 518
microsatellite genotyping, 518
natural product, 1746
ophiostomatoid fungi, 1640
southern yellow pine, 1640
survival, 518
termiticide, 1746
tetrahydrocarvone, 1746
Frankliniella bispinosa,
economic injury level, 480
flower thrips, 480
southern highbush blueberry, 480
Vaccinium corymbosum × *Vaccinium darrowi*, 480
Frankliniella fusca,
imidacloprid, 884
Neohydatothrips variabilis, 884
thiamethoxam, 884
Frankliniella occidentalis,
coating, 1238
essential oil, 1238
polymer, 1238
thrip, 1238
fruit fly, 371, 1557, 1940
biodemographic study, 371
Ceratitis rosa, 371
detection, 1557
eradication program, 1557
hot water, 1940
hydrocooling, 1940
longevity, 371
mango fruit, 1940
survey, 1557
Tephritidae, 371
thermal sensitivity, 1940
fruit fly management,
ammonium acetate, 471
Ceratitis capitata, 471
spinosad, 471
trimethylamine, 471
fumigant,
aluminum phosphide, 135
Lyctus africanus, 135
Mangifera indica, 135
Pinus roxburghii, 135
fumigation, 282, 686, 1107, 2187
ambulatory behavior, 2187
behavioral resistance, 2187
flour mill, 686
integrated pest management (IPM), 686
Lepinotus, 282
lesser grain borer, 1107
Liposcelis, 282
ozone susceptibility, 2187
phosphine resistance, 1107
Psocoptera, 282
red flour beetle, 1107
spatial distribution, 686
stored product protection, 2187
stored-product, 1107
sulfuryl fluoride, 282
Tribolium castaneum, 686
fuscumol acetate,
Cerambycidae, 906
2,3-hexanediol, 906
3-hydroxy-2-hexanone, 906
inhibition, 906

G

GABA receptor,
fipronil resistance, 1781
Oulema oryzae Kuwayama, 1781
rice leaf beetle, 1781
gall midge,
Mayetiola destructor, 1988
seasonal abundance, 1988
sex pheromone, 1988
Triticum aestivum, 1988
gene pyramid,
Aphis glycines, 1835
host plant resistance, 1835
soybean, 1835
general esterase,
acetylcholinesterase insensitivity, 540
carbamate, 540
glutathione-*S*-transferase, 540
huanglongbing, 540
generalized linear mixed model,
Bartlett's sequential test, 1077
binomial sampling, 1077
genetic differentiation,
Aphis spiraecola, 583
genetic diversity, 583
microsatellite, 583
genetic diversity, 583, 1057
adaptation, 1057
Aphis spiraecola, 583
Diuraphis noxia, 1057
genetic differentiation, 583
microsatellite, 583
plant resistance, 1057
wheat, 1057
genetic drift,
Bactrocera tryoni, 1051
domestication, 1051
mass rearing, 1051
selection, 1051
genetic group,
Bemisia tabaci, 242
diagnostic technique, 242
mitochondrial cytochrome oxidase I gene, 242
polymerase chain reaction-restriction fragment length polymorphism, 242

genetic resistance,
Apis mellifera, 313
crop pollination, 313
migratory beekeeping, 313
Varroa destructor, 313
genetically engineered crop,
Bacillus thuringiensis (*Bt*), 767
Diabrotica, 767
refuge, 767
resistance, 767
genetically modified crop,
entomopathogenic nematode, 994
meta-analysis, 994
Pectinophora gossypiella, 994
tritrophic interaction, 994
genotype,
aphid management strategy, 608
Aphis glycines, 608
Glycine max, 608
host plant resistance, 608
geographic virus isolate,
baculovirus, 344
Lymantria dispar asiatica, 344
pathogenicity, 344
virulence, 344
GF-120,
citrus orchard, 1277
hydrolyzed-protein bait, 1277
monitoring, 1277
phenology, 1277
glassy-winged sharpshooter,
biological control, 1532
egg parasitoid, 1532
parasitoid biology, 1532
Glossinidae,
Africa, 1477
insect control, 1477
livestock disease, 1477
simulation modeling, 1477
glutathione transferase,
carboxylesterase, 1034
cytochrome P450, 1034
quercetin, 1034
silkworm, 1034
glutathione-S-transferase,
acetylcholinesterase insensitivity, 540
carbamate, 540
general esterase, 540
huanglongbing, 540
Glycine max, 608, 1629
aphid management strategy, 608
Aphis glycines, 608
Armadillidium vulgare, 1629
genotype, 608
host plant resistance, 608
isopod, 1629
residue management, 1629
soybean, 1629
granular bait,
Asian needle ant, 1751
control, 1751
hydramethylnon, 1751
Pachycondyla chinensis, 1751
grape,
integrated pest management (IPM), 936
Isomate, 936
Lepidoptera, 936
semiochemical, 936
grape mealybug,
pheromone, 949
Pseudococcus maritimus, 949
reproductive biology, 949
sexual reproduction, 949
Grapholita molesta, 1426, 1712
apple, 1712
bioassay, 1426
insecticide resistance, 1426
integrated pest management (IPM), 1712
joint action, 1426
oriental fruit moth, 1712
peach, 1712
grasshopper,
application, 1915
IGR, 1915
microsporidia, 1915
green leaf volatile,
electroantennogram, 1149
Plutella xylostella, 1149
sex pheromone, 1149
trap catch, 1149
greenhouse, 34, 505, 1871
Bombus ephippiatus, 1871
bumble bee, 34, 1871
floriculture, 505
foraging tradeoff, 34
light intensity, 505
pest management, 505
pollination, 34, 1871
systemic insecticide, 505
tomato, 34, 1871
grower recommendation,
crucifer, 1719
field predation, 1719
spined soldier bug, 1719
yellowmargined leaf beetle, 1719
Gryon japonicum,
adult longevity, 1524
emergence, 1524
Ooencyrtus nezarae, 1524
parasitism, 1524
guttation,
ELISA, 67
imidacloprid, 67
melon, 67

H

haplotype,
cytochrome oxidase I, 418
Spodoptera frugiperda, 418
triose phosphate isomerase, 418
hatching,
development, 964
eclosion, 964
mating, 964
vegetative insecticidal protein, 964
heat shock,
apple, 1540
Carposina sasakii, 1540
CATTS, 1540
heat treatment, 703, 709
economics, 709
efficacy assessment, 703
insect mortality, 709
red flour beetle, 703
sanitation, 703, 709
temperature, 703, 709
Helicoverpa armigera, 1439, 2172
antibiotic resistance, 2172
canola cultivar, 2172
demographic parameter, 2172
digestive enzyme, 1439
feeding performance, 1439
host plant, 1439
life table, 2172
Helicoverpa zea,
Bacillus thuringiensis (*Bt*), 1773
Ostrinia nubilalis, 1773
refuge, 1773
Hemiptera, 616, 1509
Bacillus thuringiensis (*Bt*), 616
biological control, 1509
Coccinellidae, 1509
Lygus, 616
predator, 1509
tarnished plant bug, 616
transgenic cotton, 616
herbivory,
biofuel feedstock, 878
fertilization, 878
Schistocerca americana, 878
Spodoptera frugiperda, 878
Hermetia illucens,
black soldier fly, 374
fermentation, 374
leachate processing, 374
Herpetogramma phaeopteralis,
degree-day, 1573
developmental rate, 1573
nonlinear model, 1573
turfgrass, 1573
Hessian fly,
resistance, 642
susceptibility, 642

systemic induced susceptibility, 642
wheat, 642
2,3-hexanediol,
Cerambycidae, 906
fuscumol acetate, 906
3-hydroxy-2-hexanone, 906
inhibition, 906
highly pathogenic avian influenza,
blow fly, 1796
Calliphora nigribarbis, 1796
insecticide-treated netting screen, 1796
permethrin, 1796
honey,
brood pheromone, 304
honey bee, 304
SuperBoost, 304
survival, 304
honey bee, 20, 304, 1130, 1134, 1145, 1487, 1878
AFB, 1487
Apis mellifera, 1145, 1487, 1878
Ascosphaera apis, 1878
Bombus, 1130
brood pheromone, 304, 1134
chalkbrood, 1878
colony growth, 1134
fatty acid, 1145
flax oil, 1145
honey, 304
insemination success, 20
lysozyme, 1878
mating number, 20
matrix, 1487
Megachile, 1130
nestmate recognition, 1145
non-*Apis* bee, 1130
nutrition, 1134
Osmia, 1130
polymerase chain reaction (PCR), 1487
protein supplement, 1134
queen, 20
SuperBoost, 304
survival, 304
honey harvesting,
Apis mellifera, 777
comb orientation, 777
nest pattern, 777
traditional hive, 777
horizontal transfer,
chlorantraniliprole, 1736
eastern subterranean termite, 1736
insecticide, 1736
Reticulitermes flavipes, 1736
host discrimination,
African fruit fly, 2068
Ceratitis cosyra, 2068
fecal matter, 2068
host-marking pheromone, 2068
host genotype,
biological control, 354
diamondback moth, 354
insecticide, 354
resistance, 354
host-marking pheromone,
African fruit fly, 2068
Ceratitis cosyra, 2068
fecal matter, 2068
host discrimination, 2068
host plant, 930, 1439, 1597
areawide, 1597
Bemisia tabaci biotype B, 930
biological invasion, 930
biology, 930
biotype Q, 930
digestive enzyme, 1439
ecology, 1597
Eoreuma, 1597
feeding performance, 1439
Helicoverpa armigera, 1439
Saccharum, 1597
host plant resistance, 25, 252, 608, 1178, 1447, 1809, 1816, 1835
antibiosis, 252
aphid management strategy, 608
Aphis glycines, 25, 608, 1835
avocado, 1178
Begomovirus, 1447
biological parameter, 1809
biotype, 1816
Diabrotica virgifera virgifera, 25
fecundity, 1809
gene pyramid, 1835
genotype, 608
Glycine max, 608
integrated pest management (IPM), 252
Mediterranean fruit fly, 1178
potato cultivar, 1809
potato tuberworm, 1809
quarantine security, 1178
Rag genes, 25
risk analysis, 1178
rotation resistance, 25
soybean, 1835
thrip *Frankliniella occidentalis*, 1816
transmission, 1447
vector, 1447
host race,
cytochrome oxidase I, 573
fall armyworm, 573
PCR-restriction fragment-length polymorphism, 573
South America, 573
host status,
Mangifera indica, 1954
mango, 1954
Thaumatotibia leucotreta, 1954
hot water,
fruit fly, 1940
hydrocooling, 1940
mango fruit, 1940
thermal sensitivity, 1940
housekeeping gene,
Aphis glycines, 1432
normalization, 1432
quantitative real-time polymerase chain reaction, 1432
huanglongbing,
acetylcholinesterase insensitivity, 540
carbamate, 540
general esterase, 540
glutathione-S-transferase, 540
human-mediated dispersal,
biological invasion, 438
firewood, 438
invasive forest pest, 438
long-distance dispersal, 438
humidity,
adult survival, 186
Bactrocera oleae, 186
reproductive biology, 186
temperature, 186
hunting billbug,
Sphenophorus venatus vestitus, 2045
turfgrass, 2045
Hurricane Katrina,
Formosan subterranean termite, 518
inundation, 518
microsatellite genotyping, 518
survival, 518
Hyadaphis foeniculi,
cotton with colored fibers, 113
damage, 113
fennel, 113
intercropping, 113
hydramethylnon,
Asian needle ant, 1751
control, 1751
granular bait, 1751
Pachycondyla chinensis, 1751
hydrocooling,
fruit fly, 1940
hot water, 1940
mango fruit, 1940
thermal sensitivity, 1940
hydrolyzed-protein bait,
citrus orchard, 1277
GF-120, 1277
monitoring, 1277
phenology, 1277

3-hydroxy-2-hexanone,
Cerambycidae, 906
fuscumol acetate, 906
2,3-hexanediol, 906
inhibition, 906
hypoxia,
biodegradation, 62
escape behavior, 62
pig manure, 62
separation, 62

I

identification,
biological control, 1520
egg structure, 1520
operculum, 1520
Orius, 1520
IGR, 1915, 2097
application, 1915
Beauveria bassiana, 2097
botanical, 2097
grasshopper, 1915
insecticide, 2097
microsporidia, 1915
survival, 2097
imidacloprid, 67, 884, 1392
discriminating dose, 1392
ELISA, 67
Frankliniella fusca, 884
guttation, 67
immature, 1392
melon, 67
Neohydatothrips variabilis, 884
Planococcus ficus, 1392
resistance monitoring, 1392
thiamethoxam, 884
immature,
discriminating dose, 1392
imidacloprid, 1392
Planococcus ficus, 1392
resistance monitoring, 1392
immigration dynamic,
Candidatus Phytoplasma mali, 322
overwintered adult, 322
psyllid vector, 322
temperature, 322
indoxacarb,
boric acid, 1139
fipronil, 1139
survival, 1139
toxic bait efficacy, 1139
induced resistance,
Atherigona soccata, 1069
biochemical mechanisms of resistance, 1069
flavonoids, 1069
Sorghum bicolor, 1069
inhibition,
Cerambycidae, 906
fuscumol acetate, 906
2,3-hexanediol, 906
3-hydroxy-2-hexanone, 906
insect behavior,
Diaphorina citri, 1492
electrical penetration graph, 1492
feeding disruption, 1492
insecticide, 1492
insect control,
Africa, 1477
Glossinidae, 1477
livestock disease, 1477
simulation modeling, 1477
insect growth regulator,
larval development, 726
Stomoxys calcitrans, 726
insect histopathology,
Bacillus thuringiensis (*Bt*), 1164
camptothecin (CPT), 1164
nucleopolyhedrovirus (NPV), 1164
synergism, 1164
insect mortality,
economics, 709
heat treatment, 709
sanitation, 709
temperature, 709
insect vector,
areawide pest management, 1268
Bactericera cockerelli, 1268
Liberibacter, 1268
zebra chip, 1268
insecticidal potential,
diatomaceous earth, 670
natural zeolite, 670
Sitophilus oryzae, 670
Tribolium castaneum, 670
insecticide, 207, 354, 410, 971, 1157, 1492, 1736, 2097, 2129
Amyelois transitella, 410
Beauveria bassiana, 2097
biological control, 354
botanical, 2097
C. quinquefasciatus, 971
chlorantraniliprole, 1736
cytochrome P450 monooxygenase, 410
diamondback moth, 354
Diaphorina citri, 1492
eastern subterranean termite, 1736
electrical penetration graph, 1492
esterases resistance, 971
feeding disruption, 1492
horizontal transfer, 1736
host genotype, 354
IGR, 2097
insect behavior, 1492
Laodelphax striatellus, 2129
Leptinotarsa decemlineata, 207
life-table parameter, 1157
myristicin, 410
naled, 971
potato, 207
resistance, 354
resistance management, 207
resistance monitoring, 2129
resistance risk assessment, 2129
resmethrin, 971
Reticulitermes flavipes, 1736
sublethal effect, 1157
survival, 2097
synergist, 410
toxicity, 1157
Trichogramma chilonis, 1157
insecticide efficacy,
lethality index, 1726
pesticide, 1726
susceptibility, 1726
toxicity, 1726
insecticide induction,
cytochrome P450 monooxygenase, 560
psocid, 560
quantitative polymerase chain reaction, 560
insecticide resistance, 872, 1006, 1426
acrinathrin, 1006
bioassay, 1426
codling moth, 872
cytochrome P450, 1006
fitness trade-off, 872
Grapholita molesta, 1426
joint action, 1426
over-expression, 1006
western flower thrip, 1006
insecticide susceptibility,
acetamiprid, 988
avermectin, 988
Brontispa longissimi, 988
β-cypermethrin, 988
insecticide-treated netting screen,
blow fly, 1796
Calliphora nigribarbis, 1796
highly pathogenic avian influenza, 1796
permethrin, 1796
insemination success,
honey bee, 20
mating number, 20
queen, 20
integrated pest management (IPM), 102, 252, 686, 739, 936, 1115, 1224, 1335, 1548, 1712
alfalfa weevil, 102
antibiosis, 252
apple, 1712
attractant, 1335
attracticidal sphere, 1548

behavioral control, 1548
bioclimatic modeling, 739, 1115
cost of management, 1115
degree-day, 102
diamondback moth, 1115
flour mill, 686
fumigation, 686
grape, 936
Grapholita molesta, 1712
host plant resistance, 252
Isomate, 936
kairomone, 1335
leaf-cutting ants, 1224
Lepidoptera, 936
Leucocoprinus gongylophorus (Heim) Moeller, 1224
monitoring, 1335
MTT, 1224
natural fungicides, 1224
oriental fruit moth, 1712
oviposition, 1335
peach, 1712
pest control, 1115
Plutella xylostella, 739
population outbreak, 739
resistance, 739
Rhagoletis, 1548
semiochemical, 936
sheep, 102
spatial distribution, 686
spinosad, 1548
targeted grazing, 102
Tribolium castaneum, 686

intercropping, 113, 854
community index, 854
cotton with colored fibers, 113
damage, 113
fennel, 113
Hyadaphis foeniculi, 113
natural enemy, 854
Sitobion avenae, 854
wheat, 854

interference,
aphid, 783
Potyvirus-transmission, 783
specificity, 783

interference competition,
biological control, 386
Pseudacteon tricuspis, 386
sex ratio, 386
social form, 386

intraplant movement,
feeding preference, 847
pest management, 847
sampling, 847
stink bug, 847

inundation,
Formosan subterranean termite, 518
Hurricane Katrina, 518
microsatellite genotyping, 518
survival, 518

invasive ant,
irradiation, 817
little fire ant, 817
phytosanitary treatment, 817
quarantine pest, 817

invasive forest pest,
biological invasion, 438
firewood, 438
human-mediated dispersal, 438
long-distance dispersal, 438

invasive fruit fly,
male lure, 1186
surveillance, 1186
Tephritidae, 1186
trapping, 1186

invasive insect,
Agrilus planipennis, 1929
ash tree, 1929
autodissemination, 1929
entomopathogenic fungus, 1929

invasive mealybug,
molecular marker, 2052
pheromone, 2052
Planococcus minor, 2052

invasive species, 1, 272, 497, 1971, 2035
Allee effect, 1
armored scale, 497
avocado, 497, 2035
biological invasion, 1
delimitation, 272
density dependence, 1
detection, 272, 497
developmental biology, 2035
Diaspididae, 2035
emerald ash borer, 272
eradication, 1
Lepidoptera, 1971
phytosanitary treatment, 1971
regulatory pest, 1971
surveillance, 272, 497
x-ray radiation, 1971

Ipomoea,
Cylas formicarius, 651
Diabrotica, 651
plant resistance, 651
white grub, 651

Ips,
aggregation pheromone, 923
reproductive isolation, 923
Scolytidae, 923

irradiation, 54, 817
invasive ant, 817
light brown apple moth, 54
little fire ant, 817
mating disruption, 54
phytosanitary treatment, 817
quarantine pest, 817
sterile insect technique, 54

irrigation system,
Empoasca vitis, 176
vineyard, 176
water stress, 176

Isomate,
grape, 936
integrated pest management (IPM), 936
Lepidoptera, 936
semiochemical, 936

isopod,
Armadillidium vulgare, 1629
Glycine max, 1629
residue management, 1629
soybean, 1629

isopropanol,
Coleoptera, 2076
kairomone, 2076
Scarabaeidae, 2076

Ixodes scapularis,
blacklegged tick, 222
entomopathogenic fungi, 222
Metarhizium anisopliae, 222
tick control, 222

Ixodiphagus,
Amblyomma, 1979
biological control, 1979
Rhipicephalus sanguineus, 1979
tick, 1979

J

joint action,
bioassay, 1426
Grapholita molesta, 1426
insecticide resistance, 1426

jute bag,
aluminum sheet bin, 1095
Callosobruchus chinensis, 1095
Cicer arietinum, 1095
natural product, 1095

K

kairomone, 169, 1335, 2076
attractant, 1335
Coleoptera, 2076
integrated pest management (IPM), 1335
isopropanol, 2076
monitoring, 1335
Otiorhynchus, 169
oviposition, 1335
plant odor, 169
Scarabaeidae, 2076
strawberry, 169
weevil, 169

L

Laodelphax striatellus, 129, 2129
Cicadulina bipunctata
Forage grass, 129
insecticide, 2129
N-formylloline, 129

resistance monitoring, 2129
resistance risk assessment, 2129
Sogatella furcifera, 129
larval development,
insect growth regulator, 726
Stomoxys calcitrans, 726
larval mortality,
Bacillus thuringiensis (*Bt*), 732
Stomoxys calcitrans, 732
temperature, 732
laurel wilt, 461, 659
attraction, 461
avocado, 659
α-copaene, 659
emergence, 461
exotic species, 461
Redbay ambrosia beetle, 659
Scolytinae, 461
sesquiterpene, 659
leachate processing,
black soldier fly, 374
fermentation, 374
Hermetia illucens, 374
leaf color,
Allium cepa, 632
olfactometer, 632
onion resistance, 632
Thrips tabaci, 632
leaf-cutting ants,
integrated pest management (IPM), 1224
Leucocoprinus gongylophorus (Heim) Moeller, 1224
MTT, 1224
natural fungicides, 1224
leafhopper,
regulated deficit irrigation (RDI), 1293
vineyard, 1293
water, 1293
leafminer,
biological control, 161
ectoparasitoid, 161
sex ratio, 161
tritrophic interactions, 161
lemon-scented tea tree oil (LSO),
deterrent, 1379
diamondback moth, 1379
lemon-scented tea tree oil (LSO), 1379
repellant, 1379
Lepidoptera, 936, 1971
grape, 936
integrated pest management (IPM), 936
invasive species, 1971
Isomate, 936
phytosanitary treatment, 1971
regulatory pest, 1971
semiochemical, 936
x-ray radiation, 1971
Lepinotus,
fumigation, 282
Liposcelis, 282
Psocoptera, 282
sulfuryl fluoride, 282
Leptinotarsa decemlineata, 207, 549
chlorantraniliprole, 549
insecticide, 207
potato, 207
resistance management, 207
synergist, 549
toxicity, 549
lesser grain borer,
fumigation, 1107
phosphine resistance, 1107
red flour beetle, 1107
stored-product, 1107
lesser mealworm,
aggregation, 2196
pheromone, 2196
pitfall, 2196
poultry, 2196
50% lethal dose,
baseline data, 1024
basic model, 1024
proportionality, 1024
weight study, 1024
lethality index,
insecticide efficacy, 1726
pesticide, 1726
susceptibility, 1726
toxicity, 1726
lettuce,
Nasonovia ribisnigri, 810
phosphine fumigation, 810
phytosanitary, 810
postharvest pest control, 810
Leucocoprinus gongylophorus (Heim) Moeller,
integrated pest management (IPM), 1224
leaf-cutting ants, 1224
MTT, 1224
natural fungicides, 1224
Liberibacter,
areawide pest management, 1268
Bactericera cockerelli, 1268
insect vector, 1268
zebra chip, 1268
life cycle,
citrus, 338
net reproductive rate, 338
parasitism, 338
life table, 402, 2172
antibiotic resistance, 2172
avocado, 402
canola cultivar, 2172
demographic parameter, 2172
Helicoverpa armigera, 2172
Perú, 402
sex pheromone, 402
Stenoma catenifer, 402
life-table parameter,
insecticide, 1157
sublethal effect, 1157
toxicity, 1157
Trichogramma chilonis, 1157
light brown apple moth, 54, 1694
disruption, 1694
Epiphyas postvittana, 1694
E11-tetradecenyl acetate, 1694
irradiation, 54
mating disruption, 54
pheromone, 1694
sterile insect technique, 54
light intensity,
floriculture, 505
greenhouse, 505
pest management, 505
systemic insecticide, 505
light trap,
date palm pest, 2062
Oryctes agamemnon arabicus, 2062
photoperiod, 2062
rhinoceros beetle, 2062
light wavelength,
alate, 2213
drywood termite, 2213
phototropic, 2213
Liposcelis,
fumigation, 282
Lepinotus, 282
Psocoptera, 282
sulfuryl fluoride, 282
little fire ant,
invasive ant, 817
irradiation, 817
phytosanitary treatment, 817
quarantine pest, 817
livestock disease,
Africa, 1477
Glossinidae, 1477
insect control, 1477
simulation modeling, 1477
long-distance dispersal,
biological invasion, 438
firewood, 438
human-mediated dispersal, 438
invasive forest pest, 438
longevity, 371, 2015
Acer platanoides, 2015
Anoplophora glabripennis, 2015
biodemographic study, 371
Ceratitis rosa, 371
fecundity, 2015
fruit fly, 371
neonicotinoid, 2015
Tephritidae, 371

low pressure,
diapausing larvae, 679
methyl bromide alternative, 679
monoterpenoid, 679
Plodia interpunctella, 679
Lyctus africanus,
aluminum phosphide, 135
fumigant, 135
Mangifera indica, 135
Pinus roxburghii, 135
Lygus, 616, 2007
Bacillus thuringiensis (*Bt*), 616
cotton, 2007
Hemiptera, 616
sampling, 2007
tarnished plant bug, 616, 2007
threshold, 2007
transgenic cotton, 616
Lymantria dispar asiatica,
baculovirus, 344
geographic virus isolate, 344
pathogenicity, 344
virulence, 344
lysozyme,
Apis mellifera, 1878
Ascosphaera apis, 1878
chalkbrood, 1878
honey bee, 1878

M

M. bicoloratus,
baculovirus, 1503
parasitoid, 1503
S. exigua, 1503
S. exigua multiple nucleopolyhedrovirus, 1503
maize, 120, 1248
Bt maize, 1248
Diabrotica virgifera virgifera, 1248
(E)-β-caryophyllene, 120
fall armyworm, 120
nontarget, 1248
pyramid, 1248
resistance, 120
volatile, 120
maize cob,
baiting system, 1758
cardboard, 1758
Coptotermes formosanus, 1758
pine wood, 1758
maize weevil,
common smut, 1457
corn earworm, 1457
principal components analysis, 1457
stink bug–discolored kernel, 1457
male lure,
invasive fruit fly, 1186
surveillance, 1186
Tephritidae, 1186
trapping, 1186
management,
bed bug, 2107
Cimex lectularius, 2107
repellency, 2107
ultrasound, 2107
Mangifera indica, 135, 1954
aluminum phosphide, 135
fumigant, 135
host status, 1954
Lyctus africanus, 135
mango, 1954
Pinus roxburghii, 135
Thaumatotibia leucotreta, 1954
mango,
host status, 1954
Mangifera indica, 1954
Thaumatotibia leucotreta, 1954
mango fruit,
fruit fly, 1940
hot water, 1940
hydrocooling, 1940
thermal sensitivity, 1940
manuka oil,
Agrilus planipennis, 429
dark green trap, 429
phoebe oil, 429
(3Z)-hexenol, 429
mass attack,
attack phase, 140
mass trapping, 140
semiochemical release rate, 140
synthetic lure, 140
mass rearing, 802, 1051
artificial diet, 802
Bactrocera tryoni, 1051
biological control, 802
Coleoptera, 802
domestication, 1051
genetic drift, 1051
parasitoid wasp, 802
selection, 1051
mass trapping,
attack phase, 140
mass attack, 140
semiochemical release rate, 140
synthetic lure, 140
mating,
development, 964
eclosion, 964
hatching, 964
vegetative insecticidal protein, 964
mating disruption,
irradiation, 54
light brown apple moth, 54
sterile insect technique, 54
mating number,
honey bee, 20
insemination success, 20
queen, 20
matrix,
AFB, 1487
Apis mellifera, 1487
honey bee, 1487
polymerase chain reaction (PCR), 1487
Mayetiola destructor,
gall midge, 1988
seasonal abundance, 1988
sex pheromone, 1988
Triticum aestivum, 1988
MCH,
Douglas-fir beetle, 451
3-methylcylcohex-2-en-1-one, 451
pheromone plume, 451
Scolytinae, 451
mealybug,
Ferrisia, 1681
pistachio, 1681
Pistacia, 1681
Pseudococcidae, 1681
Mediterranean fruit fly,
avocado, 1178
host plant resistance, 1178
quarantine security, 1178
risk analysis, 1178
Megachile,
Bombus, 1130
honey bee, 1130
non-*Apis* bee, 1130
Osmia, 1130
meliponini,
biorational, 1234
pesticide, 1234
Saccharopolyspora spinosa, 1234
social, 1234
melon,
ELISA, 67
guttation, 67
imidacloprid, 67
meta-analysis,
entomopathogenic nematode, 994
genetically modified crop, 994
Pectinophora gossypiella, 994
tritrophic interaction, 994
Metarhizium anisopliae,
blacklegged tick, 222
entomopathogenic fungi, 222
Ixodes scapularis, 222
tick control, 222
method,
Cocos nucifera, 1419
pesticide, 1419
standardization, 1419

methoprene,
concrete, 718
residual efficacy, 718
surface treatment, 718
varnished wood, 718
methoxyfenozide, 1702, 1982
chrysopid, 1982
cypermethrin, 1982
endosulfan, 1982
navel orangeworm, 1702
obliquebanded leafroller, 1702
oriental fruit moth, 1702
peach twig borer, 1702
spinosad, 1982
methyl bromide alternative, 679, 1171
controlled atmosphere, 1171
diapausing larvae, 679
disinfestation, 1171
ethanol, 1171
low pressure, 679
modified atmosphere, 1171
monoterpenoid, 679
Plodia interpunctella, 679
3-methylcylcohex-2-en-1-one,
Douglas-fir beetle, 451
MCH, 451
pheromone plume, 451
Scolytinae, 451
Mexican fruit fly,
Citrus limon, 363
Rutaceae, 363
Tephritidae, 363
microsatellite, 523, 583, 2136
Aphis spiraecola, 583
colony elimination, 523
colony suppression, 523
cotton, 2136
genetic differentiation, 583
genetic diversity, 583
pest management, 2136
resistance, 2136
Reticulitermes flavipes, 523
Reticulitermes hageni, 523
soybean, 2136
microsatellite genotyping,
Formosan subterranean termite, 518
Hurricane Katrina, 518
inundation, 518
survival, 518
microsatellite marker,
Bemisia argentifolii, 753
mitochondrial cytochrome oxidase I subunit, 753
New World biotype, 753
microsporidia,
application, 1915
grasshopper, 1915
IGR, 1915
migration,
dispersal, 259
forecasting, 259
migratory beekeeping,
Apis mellifera, 313
crop pollination, 313
genetic resistance, 313
Varroa destructor, 313
mirid,
Apolygus lucorum, 1921
artificial diet, 1921
rearing, 1921
successive generations, 1921
miticide,
Apis mellifera, 1895
pesticide, 1895
queen, 1895
Varroa destructor, 1895
mitochondrial cytochrome oxidase I,
phylogenetic analysis, 48
whitefly, 48
mitochondrial cytochrome oxidase I gene,
Bemisia tabaci, 242
diagnostic technique, 242
genetic group, 242
polymerase chain reaction-restriction fragment length polymorphism, 242
mitochondrial cytochrome oxidase I subunit,
Bemisia argentifolii, 753
microsatellite marker, 753
New World biotype, 753
mitochondrial DNA, 555, 1043, 2147
Anastrepha obliqua, 2147
COI, 2147
morphology, 1043
Oligonychus, 1043
PCR-SSP, 555
phylogenetic relationship, 1043
phylogeny, 555
rDNA, 1043
reproductive form, 555
sequencing, 2147
Thrips tabaci, 555
West Indian fruit fly, 2147
mixture toxicity,
antagonism, 85
cumulative exposure, 85
ecotoxicology, 85
pesticide, 85
modified atmosphere,
controlled atmosphere, 1171
disinfestation, 1171
ethanol, 1171
methyl bromide alternative, 1171
molecular marker,
invasive mealybug, 2052
pheromone, 2052
Planococcus minor, 2052
monitoring, 837, 1277, 1335, 1350
attractant, 1335
Cerambycidae, 837
citrus orchard, 1277
detection, 1350
Drosophila suzukii, 1350
flight-intercept trap, 837
GF-120, 1277
hydrolyzed-protein bait, 1277
integrated pest management (IPM), 1335
kairomone, 1335
oviposition, 1335
phenology, 1277
trapping, 837, 1350
wood-borer detection, 837
monoterpenoid,
diapausing larvae, 679
low pressure, 679
methyl bromide alternative, 679
Plodia interpunctella, 679
morphology,
mitochondrial DNA, 1043
Oligonychus, 1043
phylogenetic relationship, 1043
rDNA, 1043
MTT,
integrated pest management (IPM), 1224
leaf-cutting ants, 1224
Leucocoprinus gongylophorus (Heim) Moeller, 1224
natural fungicides, 1224
multiplex PCR,
quarantine, 1100
RAPD, 1100
SCAR, 1100
stored-product insect, 1100
myristicin,
Amyelois transitella, 410
cytochrome P450 monooxygenase, 410
insecticide, 410
synergist, 410

N

naled,
C. quinquefasciatus, 971
esterases resistance, 971
insecticide, 971
resmethrin, 971
Nasonovia ribisnigri,
lettuce, 810
phosphine fumigation, 810
phytosanitary, 810
postharvest pest control, 810

native natural enemy,
biological control, 1311
exotic invasive pest, 1311
Necremnus spp., 1311
natural enemy, 490, 854
community index, 854
conservation biological control, 490
intercropping, 854
potato psyllid, 490
Sitobion avenae, 854
tomato psyllid, 490
toxicity, 490
wheat, 854
natural fumigant,
botanical insecticide, 1329
Camptomyia corticalis, 1329
cecidomyiid gall midge, 1329
essential oil, 1329
natural fungicides,
integrated pest management (IPM), 1224
leaf-cutting ants, 1224
Leucocoprinus gongylophorus (Heim) Moeller, 1224
MTT, 1224
natural product, 1095, 1746
aluminum sheet bin, 1095
Callosobruchus chinensis, 1095
Cicer arietinum, 1095
Formosan subterranean termite, 1746
jute bag, 1095
termiticide, 1746
tetrahydrocarvone, 1746
natural zeolite,
diatomaceous earth, 670
insecticidal potential, 670
Sitophilus oryzae, 670
Tribolium castaneum, 670
navel orangeworm,
methoxyfenozide, 1702
obliquebanded leafroller, 1702
oriental fruit moth, 1702
peach twig borer, 1702
neckband,
amitraz, 2207
collar, 2207
tick control, 2207
white-tailed deer, 2207
Necremnus spp.,
biological control, 1311
exotic invasive pest, 1311
native natural enemy, 1311
Neohydatothrips variabilis,
Frankliniella fusca, 884
imidacloprid, 884
thiamethoxam, 884
neonate,
chemical control, 1998
Eoreuma loftini, 1998
novaluron, 1998
sugarcane, 1998
neonicotinoid,
Acer platanoides, 2015
Anoplophora glabripennis, 2015
fecundity, 2015
longevity, 2015
Nesidiocoris tenuis,
cultivar, 2085
release strategy, 2085
Trichogramma achaeae, 2085
Tuta absoluta, 2085
nest pattern,
Apis mellifera, 777
comb orientation, 777
honey harvesting, 777
traditional hive, 777
nestmate recognition,
Apis mellifera, 1145
fatty acid, 1145
flax oil, 1145
honey bee, 1145
net reproductive rate,
citrus, 338
life cycle, 338
parasitism, 338
New World biotype,
Bemisia argentifolii, 753
microsatellite marker, 753
mitochondrial cytochrome oxidase I subunit, 753
N-formylloline,
Cicadulina bipunctata
Forage grass, 129
Laodelphax striatellus, 129
Sogatella furcifera, 129
no-choice,
aphid, 1844
performance, 1844
probing, 1844
resistance, 1844
non-*Apis* bee,
Bombus, 1130
honey bee, 1130
Megachile, 1130
Osmia, 1130
nonhost angiosperm volatile, 149, 1668
acetophenone, 149, 1668
Dendroctonus ponderosae, 149
pest management, 149, 1668
Pinus ponderosa, 149, 1668
verbenone, 1668
nonlinear model,
degree-day, 1573
developmental rate, 1573
Herpetogramma phaeopteralis, 1573
turfgrass, 1573
nontarget,
Bt maize, 1248
Diabrotica virgifera virgifera, 1248
maize, 1248
pyramid, 1248
normalization,
Aphis glycines, 1432
housekeeping gene, 1432
quantitative real-time polymerase chain reaction, 1432
Norway spruce,
cone insect, 916
Dioryctria abietella, 916
Eupithecia abietaria, 916
seed insect, 916
novaluron, 866, 1998
chemical control, 1998
Chlorantraniliprole, 866
Eoreuma loftini, 1998
flubendiamide, 866
neonate, 1998
spinetoram, 866
spirotetramat, 866
sugarcane, 1998
noviflumuron bait,
colony elimination, 533
subterranean termite, 533
termite baiting, 533
nucleopolyhedrovirus (NPV),
Bacillus thuringiensis (*Bt*), 1164
camptothecin (CPT), 1164
insect histopathology, 1164
nucleopolyhedrovirus (NPV), 1164
synergism, 1164
nutrition,
brood pheromone, 1134
colony growth, 1134
honey bee, 1134
protein supplement, 1134
nutritional quality,
development, 1471
reproduction, 1471
stored product, 1471
Tribolium castaneum, 1471

O

obliquebanded leafroller,
methoxyfenozide, 1702
navel orangeworm, 1702
oriental fruit moth, 1702
peach twig borer, 1702
olfactometer,
Allium cepa, 632
leaf color, 632
onion resistance, 632
Thrips tabaci, 632
Oligonychus,
mitochondrial DNA, 1043
morphology, 1043
phylogenetic relationship, 1043
rDNA, 1043

onion resistance,
Allium cepa, 632
leaf color, 632
olfactometer, 632
Thrips tabaci, 632
onions thrips,
tobacco thrips, 40
Tospovirus, 40
Ooencyrtus nezarae,
adult longevity, 1524
emergence, 1524
Gryon japonicum, 1524
parasitism, 1524
operculum,
biological control, 1520
egg structure, 1520
identification, 1520
Orius, 1520
ophiostomatoid fungi,
bark beetle, 1640
blue-stained sapwood, 1640
Formosan subterranean termite, 1640
southern yellow pine, 1640
oral,
acute toxicity, 1890
Apis mellifera anatoliaca, 1890
flumethrin, 1890
oriental fruit moth, 1702, 1712
apple, 1712
Grapholita molesta, 1712
integrated pest management (IPM), 1712
methoxyfenozide, 1702
navel orangeworm, 1702
obliquebanded leafroller, 1702
peach, 1712
peach twig borer, 1702
orientation,
Aphelinus mali, 1342
color, 1342
trap, 1342
Orius,
biological control, 1520
egg structure, 1520
identification, 1520
operculum, 1520
Oryctes agamemnon arabicus,
date palm pest, 2062
light trap, 2062
photoperiod, 2062
rhinoceros beetle, 2062
Osmia,
Bombus, 1130
honey bee, 1130
Megachile, 1130
non-*Apis* bee, 1130
Ostrinia nubilalis,
Bacillus thuringiensis (*Bt*), 1773
Helicoverpa zea, 1773
refuge, 1773
Otiorhynchus,
kairomone, 169
plant odor, 169
strawberry, 169
weevil, 169
Oulema oryzae Kuwayama,
fipronil resistance, 1781
GABA receptor, 1781
rice leaf beetle, 1781
over-expression,
acrinathrin, 1006
cytochrome P450, 1006
insecticide resistance, 1006
western flower thrip, 1006
overwintered adult,
Candidatus Phytoplasma mali, 322
immigration dynamic, 322
psyllid vector, 322
temperature, 322
oviposition,
attractant, 1335
integrated pest management (IPM), 1335
kairomone, 1335
monitoring, 1335
oviposition puncture to larvae ratio,
plant resistance, 598
soybean petiole morphology, 598
soybean plant introduction, 598
vascular histochemistry, 598
ozone susceptibility,
ambulatory behavior, 2187
behavioral resistance, 2187
fumigation, 2187
stored product protection, 2187

P

P746,
antibiosis, 2167
aphid resistance, 2167
single dominant gene, 2167
soybean, 2167
Pachycondyla chinensis,
Asian needle ant, 1751
control, 1751
granular bait, 1751
hydramethylnon, 1751
Paenibacillus,
Cyclocephala, 1320
Popillia japonica, 1320
soil insect, 1320
Tiphiidae, 1320
pan trap,
electroantennogram, 890
plant volatile, 890
sex pheromone, 890
soybean aphid, 890
parasitism, 338, 1524
adult longevity, 1524
citrus, 338
emergence, 1524
Gryon japonicum, 1524
life cycle, 338
net reproductive rate, 338
Ooencyrtus nezarae, 1524
parasitoid, 792, 1503
Agrilus planipennis, 792
baculovirus, 1503
Eulophidae, 792
fitness, 792
M. bicoloratus, 1503
S. exigua, 1503
S. exigua multiple nucleopolyhedrovirus, 1503
Tetrastichus planipennisi, 792
parasitoid biology,
biological control, 1532
egg parasitoid, 1532
glassy-winged sharpshooter, 1532
parasitoid wasp,
artificial diet, 802
biological control, 802
Coleoptera, 802
mass rearing, 802
pathogenicity,
baculovirus, 344
geographic virus isolate, 344
Lymantria dispar asiatica, 344
virulence, 344
PCR-restriction fragment-length polymorphism,
cytochrome oxidase I, 573
fall armyworm, 573
host race, 573
South America, 573
PCR-SSP,
mitochondrial DNA, 555
phylogeny, 555
reproductive form, 555
Thrips tabaci, 555
peach,
apple, 1712
Grapholita molesta, 1712
integrated pest management (IPM), 1712
oriental fruit moth, 1712
peach twig borer,
methoxyfenozide, 1702
navel orangeworm, 1702
obliquebanded leafroller, 1702
oriental fruit moth, 1702
Pectinophora gossypiella,
entomopathogenic nematode, 994
genetically modified crop, 994
meta-analysis, 994
tritrophic interaction, 994

pepper,
pest management tactic, 943
production risk, 943
tomato, 943
tomato-spotted wilt virus, 943
performance,
aphid, 1844
no-choice, 1844
probing, 1844
resistance, 1844
permethrin,
blow fly, 1796
Calliphora nigribarbis, 1796
highly pathogenic avian influenza, 1796
insecticide-treated netting screen, 1796
Perú,
avocado, 402
life table, 402
sex pheromone, 402
Stenoma catenifer, 402
pest control,
bioclimatic modeling, 1115
cost of management, 1115
diamondback moth, 1115
integrated pest management (IPM), 1115
pest management, 149, 505, 847, 1668, 2136
acetophenone, 149, 1668
cotton, 2136
Dendroctonus ponderosae, 149
feeding preference, 847
floriculture, 505
greenhouse, 505
intraplant movement, 847
light intensity, 505
microsatellite, 2136
nonhost angiosperm volatile, 149, 1668
Pinus ponderosa, 149, 1668
resistance, 2136
sampling, 847
soybean, 2136
stink bug, 847
systemic insecticide, 505
verbenone, 1668
pest management strategy,
Apolygus lucorum, 1603
early season host plant, 1603
pest management tactic,
pepper, 943
production risk, 943
tomato, 943
tomato-spotted wilt virus, 943
pest resistance,
Diabrotica virgifera virgifera, 1407
feeding behavior, 1407
transgenic maize, 1407
pest sampling,
cotton fleahopper, 896
Creontiades signatus, 896
Pseudatomoscelis seriatus, 896
verde plant bug, 896
pesticide, 85, 1234, 1419, 1726, 1895
antagonism, 85
Apis mellifera, 1895
biorational, 1234
Cocos nucifera, 1419
cumulative exposure, 85
ecotoxicology, 85
insecticide efficacy, 1726
lethality index, 1726
meliponini, 1234
method, 1419
miticide, 1895
mixture toxicity, 85
queen, 1895
Saccharopolyspora spinosa, 1234
social, 1234
standardization, 1419
susceptibility, 1726
toxicity, 1726
Varroa destructor, 1895
phenology, 1205, 1277
citrus orchard, 1277
cryptic species, 1205
Dasineura oxycoccana, 1205
GF-120, 1277
hydrolyzed-protein bait, 1277
monitoring, 1277
Vaccinium corymbosum, 1205
Vaccinium macrocarpon, 1205
pheromone, 949, 1694, 2052, 2196
aggregation, 2196
disruption, 1694
Epiphyas postvittana, 1694
E11-tetradecenyl acetate, 1694
grape mealybug, 949
invasive mealybug, 2052
lesser mealworm, 2196
light brown apple moth, 1694
molecular marker, 2052
pitfall, 2196
Planococcus minor, 2052
poultry, 2196
Pseudococcus maritimus, 949
reproductive biology, 949
sexual reproduction, 949
pheromone plume,
Douglas-fir beetle, 451
MCH, 451
3-methylcylcohex-2-en-1-one, 451
Scolytinae, 451
pheromone trap,
Cerambycidae, 395
Fluon, 395
wood-boring insect, 395
phloem-limited bacteria,
plant disease, 329
sap-sucking insect, 329
weed, 329
phoebe oil,
Agrilus planipennis, 429
dark green trap, 429
manuka oil, 429
(3Z)-hexenol, 429
phosphine fumigation,
lettuce, 810
Nasonovia ribisnigri, 810
phytosanitary, 810
postharvest pest control, 810
phosphine resistance,
fumigation, 1107
lesser grain borer, 1107
red flour beetle, 1107
stored-product, 1107
photoperiod,
date palm pest, 2062
light trap, 2062
Oryctes agamemnon arabicus, 2062
rhinoceros beetle, 2062
phototropic,
alate, 2213
drywood termite, 2213
light wavelength, 2213
phylogenetic analysis,
mitochondrial cytochrome oxidase I, 48
whitefly, 48
phylogenetic relationship,
mitochondrial DNA, 1043
morphology, 1043
Oligonychus, 1043
rDNA, 1043
phylogeny,
mitochondrial DNA, 555
PCR-SSP, 555
reproductive form, 555
Thrips tabaci, 555
phytosanitary, 810, 1963
Bactrocera invadens, 1963
cold disinfestation, 1963
commodity treatment, 1963
lettuce, 810
Nasonovia ribisnigri, 810
phosphine fumigation, 810
postharvest pest control, 810
quarantine, 1963
phytosanitary treatment, 817, 1971
invasive ant, 817
invasive species, 1971
irradiation, 817
Lepidoptera, 1971
little fire ant, 817
quarantine pest, 817
regulatory pest, 1971
x-ray radiation, 1971

pig manure,
biodegradation, 62
escape behavior, 62
hypoxia, 62
separation, 62
pine wood,
baiting system, 1758
cardboard, 1758
Coptotermes formosanus, 1758
maize cob, 1758
Pinus ponderosa, 149, 1668
acetophenone, 149, 1668
Dendroctonus ponderosae, 149
nonhost angiosperm volatile, 149, 1668
pest management, 149, 1668
verbenone, 1668
Pinus roxburghii,
aluminum phosphide, 135
fumigant, 135
Lyctus africanus, 135
Mangifera indica, 135
pistachio,
Ferrisia, 1681
mealybug, 1681
Pistacia, 1681
Pseudococcidae, 1681
Pistacia,
Ferrisia, 1681
mealybug, 1681
pistachio, 1681
Pseudococcidae, 1681
pitfall,
aggregation, 2196
lesser mealworm, 2196
pheromone, 2196
poultry, 2196
Planococcus ficus,
discriminating dose, 1392
imidacloprid, 1392
immature, 1392
resistance monitoring, 1392
Planococcus minor,
invasive mealybug, 2052
molecular marker, 2052
pheromone, 2052
plant defense activator,
reflective mulch, 1302
thrip vector, 1302
Tospovirus, 1302
plant disease,
phloem-limited bacteria, 329
sap-sucking insect, 329
weed, 329
plant odor,
kairomone, 169
Otiorhynchus, 169
strawberry, 169
weevil, 169
plant resistance, 598, 651, 1057, 1465
adaptation, 1057
antixenosis, 1465
Cylas formicarius, 651
Diabrotica, 651
Diuraphis noxia, 1057
electrical penetration graph, 1465
feeding behavior, 1465
genetic diversity, 1057
Ipomoea, 651
oviposition puncture to larvae ratio, 598
soybean petiole morphology, 598
soybean plant introduction, 598
vascular histochemistry, 598
wheat, 1057
white grub, 651
plant resistance categories,
bird cherry-oat aphid, 592
wheat line, 592
plant volatile,
electroantennogram, 890
pan trap, 890
sex pheromone, 890
soybean aphid, 890
Plodia interpunctella,
diapausing larvae, 679
low pressure, 679
methyl bromide alternative, 679
monoterpenoid, 679
Plutella xylostella, 739, 1019, 1149, 1591
bioclimatic modeling, 739
chlorantraniliprole, 1019
electroantennogram, 1149
green leaf volatile, 1149
integrated pest management (IPM), 739
population outbreak, 739
resistance, 739
resistance monitoring, 1019
sex pheromone, 1149
Solenopsis geminata, 1591
Solenopsis invicta, 1591
trap catch, 1149
venom alkaloids, 1591
pollination, 34, 1871
Bombus ephippiatus, 1871
bumble bee, 34, 1871
foraging tradeoff, 34
greenhouse, 34, 1871
tomato, 34, 1871
pollinator,
alfalfa, 14
fluctuating thermal regime, 14
spring incubation, 14
polymer,
coating, 1238
essential oil, 1238
Frankliniella occidentalis, 1238
thrip, 1238
polymerase chain reaction (PCR),
AFB, 1487
Apis mellifera, 1487
honey bee, 1487
matrix, 1487
polymerase chain reaction (PCR), 1487
polymerase chain reaction-restriction fragment length polymorphism,
Bemisia tabaci, 242
diagnostic technique, 242
genetic group, 242
mitochondrial cytochrome oxidase I gene, 242
Popillia japonica, 1320, 1688
Cyclocephala, 1320
Paenibacillus, 1320
precipitation, 1688
soil insect, 1320
Tiphiidae, 1320
wash-off, 1688
population density,
Rhyzopertha dominica, 1194
spatial distribution, 1194
temporal continuity, 1194
population dynamics, 288, 823
Anastrepha, 823
areawide pest management, 823
climate, 823
copper, 288
development, 288
Spodoptera litura, 288
time series analysis, 823
population growth,
Aethina tumida, 26
diet, 26
resource acquisition, 26
temperature, 26
population outbreak,
bioclimatic modeling, 739
integrated pest management (IPM), 739
Plutella xylostella, 739
resistance, 739
postharvest pest control,
lettuce, 810
Nasonovia ribisnigri, 810
phosphine fumigation, 810
phytosanitary, 810
potato,
insecticide, 207
Leptinotarsa decemlineata, 207
resistance management, 207

potato cultivar,
biological parameter, 1809
fecundity, 1809
host plant resistance, 1809
potato tuberworm, 1809
potato psyllid,
conservation biological control, 490
natural enemy, 490
tomato psyllid, 490
toxicity, 490
potato tuberworm,
biological parameter, 1809
fecundity, 1809
host plant resistance, 1809
potato cultivar, 1809
potato virus Y, 1903, 1909
acquisition, 1903
aphid, 1909
aphid behavior, 1903
artificial diet, 1903
DNA barcoding, 1909
vector, 1909
Potyvirus-transmission,
aphid, 783
interference, 783
specificity, 783
poultry,
aggregation, 2196
lesser mealworm, 2196
pheromone, 2196
pitfall, 2196
precipitation,
Popillia japonica, 1688
wash-off, 1688
predator,
biological control, 1509
Coccinellidae, 1509
Hemiptera, 1509
preemptive removal,
economic analysis, 196
emerald ash borer, 196
urban forest management, 196
principal components analysis,
common smut, 1457
corn earworm, 1457
maize weevil, 1457
stink bug–discolored kernel, 1457
probing,
aphid, 1844
no-choice, 1844
performance, 1844
resistance, 1844
production risk,
pepper, 943
pest management tactic, 943
tomato, 943
tomato-spotted wilt virus, 943
proportionality,
baseline data, 1024
basic model, 1024
50% lethal dose, 1024
weight study, 1024
protein supplement,
brood pheromone, 1134
colony growth, 1134
honey bee, 1134
nutrition, 1134
Pseudacteon tricuspis,
biological control, 386
interference competition, 386
sex ratio, 386
social form, 386
Pseudatomoscelis seriatus,
cotton fleahopper, 896
Creontiades signatus, 896
pest sampling, 896
verde plant bug, 896
Pseudococcidae,
Ferrisia, 1681
mealybug, 1681
pistachio, 1681
Pistacia, 1681
Pseudococcus maritimus,
grape mealybug, 949
pheromone, 949
reproductive biology, 949
sexual reproduction, 949
psocid,
cytochrome P450 monooxygenase, 560
insecticide induction, 560
quantitative polymerase chain reaction, 560
Psocoptera,
fumigation, 282
Lepinotus, 282
Liposcelis, 282
sulfuryl fluoride, 282
psyllid vector,
Candidatus Phytoplasma mali, 322
immigration dynamic, 322
overwintered adult, 322
temperature, 322
pulegone,
citronellal, 72
eugenol, 72
flight muscles, 72
α-terpineol, 72
pyramid,
Bt maize, 1248
Diabrotica virgifera virgifera, 1248
maize, 1248
nontarget, 1248
pyrethroid, 957, 1612, 1789
bed bug, 1789
bug bomb, 957
Cimex lectularius, 957, 1789
desiccant dust, 1789
economic analysis, 1612
fogger, 957
regional survey, 1612
resistance, 957, 1789
risk aversion, 1612

Q

quantitative polymerase chain reaction,
cytochrome P450 monooxygenase, 560
insecticide induction, 560
psocid, 560
quantitative real-time polymerase chain reaction,
Aphis glycines, 1432
housekeeping gene, 1432
normalization, 1432
quarantine, 297, 1100, 1963
apple, 297
Bactrocera invadens, 1963
codling moth, 297
cold disinfestation, 1963
commodity treatment, 1963
Cydia pomonella, 297
diapause, 297
multiplex PCR, 1100
phytosanitary, 1963
RAPD, 1100
SCAR, 1100
stored-product insect, 1100
quarantine pest,
invasive ant, 817
irradiation, 817
little fire ant, 817
phytosanitary treatment, 817
quarantine security,
avocado, 1178
host plant resistance, 1178
Mediterranean fruit fly, 1178
risk analysis, 1178
queen, 20, 1895
Apis mellifera, 1895
honey bee, 20
insemination success, 20
mating number, 20
miticide, 1895
pesticide, 1895
Varroa destructor, 1895
quercetin,
carboxylesterase, 1034
cytochrome P450, 1034
glutathione transferase, 1034
silkworm, 1034

R

Rag genes,
Aphis glycines, 25
Diabrotica virgifera virgifera, 25
host plant resistance, 25
rotation resistance, 25

range estimation,
Abbott's formula, 1856
Bayesian statistics, 1856
EC_{50}, 1856
repellency, 1856
RAPD,
multiplex PCR, 1100
quarantine, 1100
SCAR, 1100
stored-product insect, 1100
rDNA,
mitochondrial DNA, 1043
morphology, 1043
Oligonychus, 1043
phylogenetic relationship, 1043
rearing,
Apolygus lucorum, 1921
artificial diet, 1921
mirid, 1921
successive generations, 1921
red flour beetle, 703, 1107
efficacy assessment, 703
fumigation, 1107
heat treatment, 703
lesser grain borer, 1107
phosphine resistance, 1107
sanitation, 703
stored-product, 1107
temperature, 703
Redbay ambrosia beetle,
avocado, 659
α-copaene, 659
laurel wilt, 659
sesquiterpene, 659
reflective mulch,
plant defense activator, 1302
thrip vector, 1302
Tospovirus, 1302
refuge, 767, 1773
Bacillus thuringiensis (*Bt*), 767, 1773
Diabrotica, 767
genetically engineered crop, 767
Helicoverpa zea, 1773
Ostrinia nubilalis, 1773
resistance, 767
regional survey,
economic analysis, 1612
pyrethroid, 1612
risk aversion, 1612
regulated deficit irrigation (RDI),
leafhopper, 1293
regulated deficit irrigation (RDI), 1293
vineyard, 1293
water, 1293
regulatory pest,
invasive species, 1971
Lepidoptera, 1971
phytosanitary treatment, 1971
x-ray radiation, 1971
release strategy,
cultivar, 2085
Nesidiocoris tenuis, 2085
Trichogramma achaeae, 2085
Tuta absoluta, 2085
repellant,
deterrent, 1379
diamondback moth, 1379
lemon-scented tea tree oil (LSO), 1379
repellency, 379, 1856, 2107
Abbott's formula, 1856
Bayesian statistics, 1856
bed bug, 2107
Cimex lectularius, 2107
EC_{50}, 1856
essential oil, 379
management, 2107
range estimation, 1856
sandalwood, 379
santalol, 379
Tetranychus urticae, 379
ultrasound, 2107
reproduction,
development, 1471
nutritional quality, 1471
stored product, 1471
Tribolium castaneum, 1471
reproductive biology, 186, 949
adult survival, 186
Bactrocera oleae, 186
grape mealybug, 949
humidity, 186
pheromone, 949
Pseudococcus maritimus, 949
sexual reproduction, 949
temperature, 186
reproductive form,
mitochondrial DNA, 555
PCR-SSP, 555
phylogeny, 555
Thrips tabaci, 555
reproductive isolation,
aggregation pheromone, 923
Ips, 923
Scolytidae, 923
residual efficacy,
concrete, 718
methoprene, 718
surface treatment, 718
varnished wood, 718
residue management,
Armadillidium vulgare, 1629
Glycine max, 1629
isopod, 1629
soybean, 1629
resistance, 120, 214, 354, 642, 739, 767, 957, 1401, 1789, 1844, 2136
acetamiprid, 1401
aphid, 1844
Bacillus thuringiensis (*Bt*), 767
baseline susceptibility, 214
bed bug, 1789
bioclimatic modeling, 739
biological control, 354
bug bomb, 957
Cimex lectularius, 957, 1789
corn borer, 214
cotton, 2136
desiccant dust, 1789
Diabrotica, 767
diamondback moth, 354
(E)-β-caryophyllene, 120
fall armyworm, 120
fitness cost, 1401
fogger, 957
genetically engineered crop, 767
Hessian fly, 642
host genotype, 354
insecticide, 354
integrated pest management (IPM), 739
maize, 120
microsatellite, 2136
no-choice, 1844
performance, 1844
pest management, 2136
Plutella xylostella, 739
population outbreak, 739
probing, 1844
pyrethroid, 957, 1789
refuge, 767
soybean, 2136
susceptibility, 642
systemic induced susceptibility, 642
transgenic maize, 214
volatile, 120
wheat, 642
whitefly, 1401
resistance management, 207, 1214
blended-seed refuge, 1214
Bt corn, 1214
dispersal, 1214
European corn borer, 1214
insecticide, 207
Leptinotarsa decemlineata, 207
potato, 207
resistance monitoring, 1019, 1392, 2129
chlorantraniliprole, 1019
discriminating dose, 1392
imidacloprid, 1392
immature, 1392
insecticide, 2129
Laodelphax striatellus, 2129
Planococcus ficus, 1392
Plutella xylostella, 1019
resistance risk assessment, 2129

resistance risk assessment,
insecticide, 2129
Laodelphax striatellus, 2129
resistance monitoring, 2129
resmethrin,
C. quinquefasciatus, 971
esterases resistance, 971
insecticide, 971
naled, 971
resource acquisition,
Aethina tumida, 26
diet, 26
population growth, 26
temperature, 26
Reticulitermes flavipes, 523, 1736
chlorantraniliprole, 1736
colony elimination, 523
colony suppression, 523
eastern subterranean termite, 1736
horizontal transfer, 1736
insecticide, 1736
microsatellite, 523
Reticulitermes hageni, 523
Reticulitermes hageni,
colony elimination, 523
colony suppression, 523
microsatellite, 523
Reticulitermes flavipes, 523
Reunion Island,
cage, 1358
spinosad, 1358
Synéis-appât, 1358
Tephritidae, 1358
Rhagoletis,
attracticidal sphere, 1548
behavioral control, 1548
integrated pest management (IPM), 1548
spinosad, 1548
rhinoceros beetle,
date palm pest, 2062
light trap, 2062
Oryctes agamemnon arabicus, 2062
photoperiod, 2062
Rhipicephalus sanguineus,
Amblyomma, 1979
biological control, 1979
Ixodiphagus, 1979
tick, 1979
Rhyzopertha dominica, 1088, 1194
population density, 1194
short exposure, 1088
Sitophilus oryzae, 1088
spatial distribution, 1194
spinetoram, 1088
temporal continuity, 1194
Tribolium confusum, 1088
rice leaf beetle,
fipronil resistance, 1781
GABA receptor, 1781
Oulema oryzae Kuwayama, 1781
risk analysis,
avocado, 1178
host plant resistance, 1178
Mediterranean fruit fly, 1178
quarantine security, 1178
risk aversion,
economic analysis, 1612
pyrethroid, 1612
regional survey, 1612
root maggot,
canola, 1261
crop rotation, 1261
Delia spp., 1261
rotation resistance,
Aphis glycines, 25
Diabrotica virgifera virgifera, 25
host plant resistance, 25
Rag genes, 25
Rutaceae,
Citrus limon, 363
Mexican fruit fly, 363
Tephritidae, 363

S

S. exigua,
baculovirus, 1503
M. bicoloratus, 1503
parasitoid, 1503
S. exigua multiple nucleopolyhedrovirus, 1503
S. exigua multiple nucleopolyhedrovirus,
baculovirus, 1503
M. bicoloratus, 1503
parasitoid, 1503
S. exigua, 1503
Saccharopolyspora spinosa,
biorational, 1234
meliponini, 1234
pesticide, 1234
social, 1234
Saccharum,
areawide, 1597
ecology, 1597
Eoreuma, 1597
host plant, 1597
Sahlbergella singularis,
cherelle wilt, 1285
cocoa mirids, 1285
cocoa yield, 1285
Theobroma cacao, 1285
salivary gland,
biotype, 232
DNA methylation, 232
sodium bisulfite, 232
virulence, 232
sampling, 847, 2007
cotton, 2007
feeding preference, 847
intraplant movement, 847
Lygus, 2007
pest management, 847
stink bug, 847
tarnished plant bug, 2007
threshold, 2007
sandalwood,
essential oil, 379
repellency, 379
santalol, 379
Tetranychus urticae, 379
sanitation, 703, 709
economics, 709
efficacy assessment, 703
heat treatment, 703, 709
insect mortality, 709
red flour beetle, 703
temperature, 703, 709
santalol,
essential oil, 379
repellency, 379
sandalwood, 379
Tetranychus urticae, 379
sap-sucking insect,
phloem-limited bacteria, 329
plant disease, 329
weed, 329
SCAR,
multiplex PCR, 1100
quarantine, 1100
RAPD, 1100
stored-product insect, 1100
Scarabaeidae,
Coleoptera, 2076
isopropanol, 2076
kairomone, 2076
Schistocerca americana,
biofuel feedstock, 878
fertilization, 878
herbivory, 878
Spodoptera frugiperda, 878
Scolytidae,
aggregation pheromone, 923
Ips, 923
reproductive isolation, 923
Scolytinae, 451, 461
attraction, 461
Douglas-fir beetle, 451
emergence, 461
exotic species, 461
laurel wilt, 461
MCH, 451
3-methylcylcohex-2-en-1-one, 451
pheromone plume, 451
seasonal abundance,
gall midge, 1988
Mayetiola destructor, 1988
sex pheromone, 1988

Triticum aestivum, 1988
seed insect,
cone insect, 916
Dioryctria abietella, 916
Eupithecia abietaria, 916
Norway spruce, 916
selection,
Bactrocera tryoni, 1051
domestication, 1051
genetic drift, 1051
mass rearing, 1051
selectivity,
Bacillus thuringiensis (*Bt*), 2122
Cnaphalocrocis medinalis, 2122
Cry toxin, 2122
susceptibility, 2122
semiochemical, 936, 2029
aggregation pheromone, 2029
grape, 936
integrated pest management (IPM), 936
Isomate, 936
Lepidoptera, 936
2-(undecyloxy)-ethanol, 2029
semiochemical release rate,
attack phase, 140
mass attack, 140
mass trapping, 140
synthetic lure, 140
separation,
biodegradation, 62
escape behavior, 62
hypoxia, 62
pig manure, 62
sequencing,
Anastrepha obliqua, 2147
COI, 2147
mitochondrial DNA, 2147
West Indian fruit fly, 2147
sesquiterpene,
avocado, 659
α-copaene, 659
laurel wilt, 659
Redbay ambrosia beetle, 659
sex pheromone, 402, 890, 1149, 1566, 1988
avocado, 402
electroantennogram, 890, 1149
gall midge, 1988
green leaf volatile, 1149
life table, 402
Mayetiola destructor, 1988
pan trap, 890
Perú, 402
plant volatile, 890
Plutella xylostella, 1149
seasonal abundance, 1988
soybean aphid, 890
Stenoma catenifer, 402
trap catch, 1149
Trichophysetis cretacea, 1566
Triticum aestivum, 1988
(Z)-11-hexadecenal, 1566
(Z)-11-hexadecenol, 1566
(Z)-11-hexadecenyl acetate, 1566
sex ratio, 161, 386, 2161
biological control, 161, 386
Cardinium, 2161
CI, 2161
ectoparasitoid, 161
fitness, 2161
interference competition, 386
leafminer, 161
Pseudacteon tricuspis, 386
social form, 386
Sogatella furcifera, 2161
tritrophic interactions, 161
sexual reproduction,
grape mealybug, 949
pheromone, 949
Pseudococcus maritimus, 949
reproductive biology, 949
sheep,
alfalfa weevil, 102
degree-day, 102
integrated pest management (IPM), 102
targeted grazing, 102
short exposure,
Rhyzopertha dominica, 1088
Sitophilus oryzae, 1088
spinetoram, 1088
Tribolium confusum, 1088
silkworm,
carboxylesterase, 1034
cytochrome P450, 1034
glutathione transferase, 1034
quercetin, 1034
silverfish,
bait, 1385
boric acid, 1385
toxicity, 1385
simulation modeling,
Africa, 1477
Glossinidae, 1477
insect control, 1477
livestock disease, 1477
single dominant gene,
antibiosis, 2167
aphid resistance, 2167
P746, 2167
soybean, 2167
Sitobion avenae,
community index, 854
intercropping, 854
natural enemy, 854
wheat, 854
Sitophilus oryzae, 670, 1088
diatomaceous earth, 670
insecticidal potential, 670
natural zeolite, 670
Rhyzopertha dominica, 1088
short exposure, 1088
spinetoram, 1088
Tribolium castaneum, 670
Tribolium confusum, 1088
social,
biorational, 1234
meliponini, 1234
pesticide, 1234
Saccharopolyspora spinosa, 1234
social form,
biological control, 386
interference competition, 386
Pseudacteon tricuspis, 386
sex ratio, 386
sodium bisulfite,
biotype, 232
DNA methylation, 232
salivary gland, 232
virulence, 232
Sogatella furcifera, 129, 2161
Cardinium, 2161
CI, 2161
Cicadulina bipunctata
fitness, 2161
Forage grass, 129
Laodelphax striatellus, 129
N-formylloline, 129
sex ratio, 2161
soil insect,
Cyclocephala, 1320
Paenibacillus, 1320
Popillia japonica, 1320
Tiphiidae, 1320
Solenopsis geminata,
Plutella xylostella, 1591
Solenopsis invicta, 1591
venom alkaloids, 1591
Solenopsis invicta,
Plutella xylostella, 1591
Solenopsis geminata, 1591
venom alkaloids, 1591
Sorghum bicolor,
Atherigona soccata, 1069
biochemical mechanisms of resistance, 1069
flavonoids, 1069
induced resistance, 1069
South America,
cytochrome oxidase I, 573
fall armyworm, 573
host race, 573
PCR-restriction fragment-length polymorphism, 573
southern highbush blueberry,
economic injury level, 480
flower thrips, 480
Frankliniella bispinosa, 480
Vaccinium corymbosum × *Vaccinium darrowi*, 480

southern yellow pine,
bark beetle, 1640
blue-stained sapwood, 1640
Formosan subterranean termite, 1640
ophiostomatoid fungi, 1640
soybean, 1629, 1835, 2136, 2167
antibiosis, 2167
aphid resistance, 2167
Aphis glycines, 1835
Armadillidium vulgare, 1629
cotton, 2136
gene pyramid, 1835
Glycine max, 1629
host plant resistance, 1835
isopod, 1629
microsatellite, 2136
P746, 2167
pest management, 2136
residue management, 1629
resistance, 2136
single dominant gene, 2167
soybean aphid,
electroantennogram, 890
pan trap, 890
plant volatile, 890
sex pheromone, 890
soybean petiole morphology,
oviposition puncture to larvae ratio, 598
plant resistance, 598
soybean plant introduction, 598
vascular histochemistry, 598
soybean plant introduction,
oviposition puncture to larvae ratio, 598
plant resistance, 598
soybean petiole morphology, 598
vascular histochemistry, 598
spatial analysis,
urban insect pest, 1659
urban landscape, 1659
spatial distribution, 686, 1194
flour mill, 686
fumigation, 686
integrated pest management (IPM), 686
population density, 1194
Rhyzopertha dominica, 1194
temporal continuity, 1194
Tribolium castaneum, 686
specificity,
aphid, 783
interference, 783
Potyvirus-transmission, 783
Sphenophorus venatus vestitus,
hunting billbug, 2045
turfgrass, 2045
spined soldier bug,
crucifer, 1719
field predation, 1719
grower recommendation, 1719
yellowmargined leaf beetle, 1719
spinetoram, 866, 1088
Chlorantraniliprole, 866
flubendiamide, 866
novaluron, 866
Rhyzopertha dominica, 1088
short exposure, 1088
Sitophilus oryzae, 1088
spirotetramat, 866
Tribolium confusum, 1088
spinosad, 471, 979, 1358, 1548, 1982
Alphitobius diaperinus, 979
ammonium acetate, 471
attracticidal sphere, 1548
behavioral control, 1548
cage, 1358
Ceratitis capitata, 471
chrysopid, 1982
cyfluthrin, 979
cypermethrin, 1982
diatomaceous earth, 979
endosulfan, 1982
field efficacy, 979
fruit fly management, 471
integrated pest management (IPM), 1548
methoxyfenozide, 1982
Reunion Island, 1358
Rhagoletis, 1548
Synéis-appât, 1358
Tephritidae, 1358
trimethylamine, 471
Spintherus dubius,
Apion spp., 1620
biological control, 1620
thiacloprid, 1620
Trifolium pratense, 1620
spirotetramat,
Chlorantraniliprole, 866
flubendiamide, 866
novaluron, 866
spinetoram, 866
Spodoptera exigua,
Diatraea grandiosella, 1825
Diatraea saccharalis, 1825
Spodoptera frugiperda, 1825
transgenic corn, 1825
Spodoptera frugiperda, 418, 878, 1825
biofuel feedstock, 878
cytochrome oxidase I, 418
Diatraea grandiosella, 1825
Diatraea saccharalis, 1825
fertilization, 878
haplotype, 418
herbivory, 878
Schistocerca americana, 878
Spodoptera exigua, 1825
transgenic corn, 1825
triose phosphate isomerase, 418
Spodoptera litura,
copper, 288
development, 288
population dynamics, 288
spring incubation,
alfalfa, 14
fluctuating thermal regime, 14
pollinator, 14
standardization,
Cocos nucifera, 1419
method, 1419
pesticide, 1419
Stenoma catenifer,
avocado, 402
life table, 402
Perú, 402
sex pheromone, 402
sterile insect technique,
irradiation, 54
light brown apple moth, 54
mating disruption, 54
stink bug,
feeding preference, 847
intraplant movement, 847
pest management, 847
sampling, 847
stink bug–discolored kernel,
common smut, 1457
corn earworm, 1457
maize weevil, 1457
principal components analysis, 1457
stink bug–discolored kernel, 1457
Stomoxys calcitrans, 726, 732, 1863
adult mortality, 1863
Bacillus thuringiensis (*Bt*), 732, 1863
diet, 1863
insect growth regulator, 726
larval development, 726
larval mortality, 732
temperature, 732
stored product, 1107, 1471
development, 1471
fumigation, 1107
lesser grain borer, 1107
nutritional quality, 1471
phosphine resistance, 1107
red flour beetle, 1107
reproduction, 1471
Tribolium castaneum, 1471
stored-product insect,
multiplex PCR, 1100
quarantine, 1100
RAPD, 1100
SCAR, 1100

stored product protection,
ambulatory behavior, 2187
behavioral resistance, 2187
fumigation, 2187
ozone susceptibility, 2187
strawberry,
kairomone, 169
Otiorhynchus, 169
plant odor, 169
weevil, 169
sublethal effect,
insecticide, 1157
life-table parameter, 1157
toxicity, 1157
Trichogramma chilonis, 1157
subterranean termite, 533, 2115
botanical insecticide, 2115
chloroform extract, 2115
colony elimination, 533
noviflumuron bait, 533
termite baiting, 533
successive generations,
Apolygus lucorum, 1921
artificial diet, 1921
mirid, 1921
rearing, 1921
sugarcane,
chemical control, 1998
Eoreuma loftini, 1998
neonate, 1998
novaluron, 1998
sulfuryl fluoride,
fumigation, 282
Lepinotus, 282
Liposcelis, 282
Psocoptera, 282
SuperBoost,
brood pheromone, 304
honey, 304
honey bee, 304
survival, 304
surface treatment,
concrete, 718
methoprene, 718
residual efficacy, 718
varnished wood, 718
surveillance, 272, 497, 1186
armored scale, 497
avocado, 497
delimitation, 272
detection, 272, 497
emerald ash borer, 272
invasive fruit fly, 1186
invasive species, 272, 497
male lure, 1186
Tephritidae, 1186
trapping, 1186
survey,
detection, 1557
eradication program, 1557
fruit fly, 1557
survival, 304, 518, 1139, 2097
Beauveria bassiana, 2097
boric acid, 1139
botanical, 2097
brood pheromone, 304
fipronil, 1139
Formosan subterranean termite, 518
honey, 304
honey bee, 304
Hurricane Katrina, 518
IGR, 2097
indoxacarb, 1139
insecticide, 2097
inundation, 518
microsatellite genotyping, 518
SuperBoost, 304
toxic bait efficacy, 1139
susceptibility, 642, 1726, 2122
Bacillus thuringiensis (*Bt*), 2122
Cnaphalocrocis medinalis, 2122
Cry toxin, 2122
Hessian fly, 642
insecticide efficacy, 1726
lethality index, 1726
pesticide, 1726
resistance, 642
selectivity, 2122
systemic induced susceptibility, 642
toxicity, 1726
wheat, 642
Synéis-appât,
cage, 1358
Reunion Island, 1358
spinosad, 1358
Tephritidae, 1358
synergism,
Bacillus thuringiensis (*Bt*), 1164
camptothecin (CPT), 1164
insect histopathology, 1164
nucleopolyhedrovirus (NPV), 1164
synergist, 410, 549
Amyelois transitella, 410
chlorantraniliprole, 549
cytochrome P450 monooxygenase, 410
insecticide, 410
Leptinotarsa decemlineata, 549
myristicin, 410
toxicity, 549
synthetic lure,
attack phase, 140
mass attack, 140
mass trapping, 140
semiochemical release rate, 140
systemic induced susceptibility,
Hessian fly, 642
resistance, 642
susceptibility, 642
wheat, 642
systemic insecticide,
floriculture, 505
greenhouse, 505
light intensity, 505
pest management, 505

T

targeted grazing,
alfalfa weevil, 102
degree-day, 102
integrated pest management (IPM), 102
sheep, 102
tarnished plant bug, 616, 2007
Bacillus thuringiensis (*Bt*), 616
cotton, 2007
Hemiptera, 616
Lygus, 616, 2007
sampling, 2007
threshold, 2007
transgenic cotton, 616
temperature, 26, 186, 322, 703, 709, 732
adult survival, 186
Aethina tumida, 26
Bacillus thuringiensis (*Bt*), 732
Bactrocera oleae, 186
Candidatus Phytoplasma mali, 322
diet, 26
economics, 709
efficacy assessment, 703
heat treatment, 703, 709
humidity, 186
immigration dynamic, 322
insect mortality, 709
larval mortality, 732
overwintered adult, 322
population growth, 26
psyllid vector, 322
red flour beetle, 703
reproductive biology, 186
resource acquisition, 26
sanitation, 703, 709
Stomoxys calcitrans, 732
temperature-dependent development,
biophysical model, 860
degree-day model, 860
western tarnished plant bug, 860
temporal continuity,
population density, 1194
Rhyzopertha dominica, 1194
spatial distribution, 1194

Tephritidae, 363, 371, 1186, 1358
biodemographic study, 371
cage, 1358
Ceratitis rosa, 371
Citrus limon, 363
fruit fly, 371
invasive fruit fly, 1186
longevity, 371
male lure, 1186
Mexican fruit fly, 363
Reunion Island, 1358
Rutaceae, 363
spinosad, 1358
surveillance, 1186
Synéis-appât, 1358
trapping, 1186
termite baiting,
colony elimination, 533
noviflumuron bait, 533
subterranean termite, 533
termite control,
Coptotermes formosanus, 1766
dust formulation, 1766
termiticide, 1766
termiticide, 1746, 1766
Coptotermes formosanus, 1766
dust formulation, 1766
Formosan subterranean termite, 1746
natural product, 1746
termite control, 1766
tetrahydrocarvone, 1746
α-terpineol,
citronellal, 72
eugenol, 72
flight muscles, 72
pulegone, 72
tetrahydrocarvone,
Formosan subterranean termite, 1746
natural product, 1746
termiticide, 1746
Tetranychus urticae,
essential oil, 379
repellency, 379
sandalwood, 379
santalol, 379
Tetrastichus planipennisi,
Agrilus planipennis, 792
Eulophidae, 792
fitness, 792
parasitoid, 792
Thaumatotibia leucotreta,
host status, 1954
Mangifera indica, 1954
mango, 1954
Theobroma cacao,
cherelle wilt, 1285
cocoa mirids, 1285
cocoa yield, 1285
Sahlbergella singularis, 1285
thermal sensitivity,
fruit fly, 1940
hot water, 1940
hydrocooling, 1940
mango fruit, 1940
thiacloprid,
Apion spp., 1620
biological control, 1620
Spintherus dubius, 1620
Trifolium pratense, 1620
thiamethoxam,
Frankliniella fusca, 884
imidacloprid, 884
Neohydatothrips variabilis, 884
threshold,
cotton, 2007
Lygus, 2007
sampling, 2007
tarnished plant bug, 2007
thrip,
coating, 1238
essential oil, 1238
Frankliniella occidentalis, 1238
polymer, 1238
thrip *Frankliniella occidentalis*,
biotype, 1816
host-plant resistance, 1816
thrip vector,
plant defense activator, 1302
reflective mulch, 1302
Tospovirus, 1302
Thrips tabaci, 555, 632
Allium cepa, 632
leaf color, 632
mitochondrial DNA, 555
olfactometer, 632
onion resistance, 632
PCR-SSP, 555
phylogeny, 555
reproductive form, 555
tick,
Amblyomma, 1979
biological control, 1979
Ixodiphagus, 1979
Rhipicephalus sanguineus, 1979
tick control, 222, 2207
amitraz, 2207
blacklegged tick, 222
collar, 2207
entomopathogenic fungi, 222
Ixodes scapularis, 222
Metarhizium anisopliae, 222
neckband, 2207
white-tailed deer, 2207
time series analysis,
Anastrepha, 823
areawide pest management, 823
climate, 823
population dynamics, 823
Tiphiidae,
Cyclocephala, 1320
Paenibacillus, 1320
Popillia japonica, 1320
soil insect, 1320
tipworm,
apical-meristem injury, 1366
cranberry, 1366
Dasineura oxycoccana, 1366
Ericaceae, 1366
tobacco thrips,
onions thrips, 40
Tospovirus, 40
tolerance,
antibiosis, 2180
antixenosis, 2180
wheat, 2180
wheat curl mite, 2180
tomato, 34, 943, 1871
Bombus ephippiatus, 1871
bumble bee, 34, 1871
foraging tradeoff, 34
greenhouse, 34, 1871
pepper, 943
pest management tactic, 943
pollination, 34, 1871
production risk, 943
tomato-spotted wilt virus, 943
tomato psyllid,
conservation biological control, 490
natural enemy, 490
potato psyllid, 490
toxicity, 490
tomato-spotted wilt virus,
pepper, 943
pest management tactic, 943
production risk, 943
tomato, 943
Tospovirus, 40, 1302
onions thrips, 40
plant defense activator, 1302
reflective mulch, 1302
thrip vector, 1302
tobacco thrips, 40
toxic bait efficacy,
boric acid, 1139
fipronil, 1139
indoxacarb, 1139
survival, 1139
toxicity, 490, 549, 1157, 1385, 1726
bait, 1385
boric acid, 1385
chlorantraniliprole, 549
conservation biological control, 490
insecticide, 1157
insecticide efficacy, 1726
Leptinotarsa decemlineata, 549
lethality index, 1726
life-table parameter, 1157
natural enemy, 490

pesticide, 1726
potato psyllid, 490
silverfish, 1385
sublethal effect, 1157
susceptibility, 1726
synergist, 549
tomato psyllid, 490
Trichogramma chilonis, 1157
traditional hive,
Apis mellifera, 777
comb orientation, 777
honey harvesting, 777
nest pattern, 777
transgenic corn,
Diatraea grandiosella, 1825
Diatraea saccharalis, 1825
Spodoptera exigua, 1825
Spodoptera frugiperda, 1825
transgenic cotton,
Bacillus thuringiensis (*Bt*), 616
Hemiptera, 616
Lygus, 616
tarnished plant bug, 616
transgenic maize, 214, 1407
Bacillus thuringiensis (*Bt*), 214
baseline susceptibility, 214
corn borer, 214
Diabrotica virgifera virgifera, 1407
feeding behavior, 1407
pest resistance, 1407
resistance, 214
transmission,
Begomovirus, 1447
host plant resistance, 1447
vector, 1447
trap,
Aphelinus mali, 1342
color, 1342
orientation, 1342
trap catch,
electroantennogram, 1149
green leaf volatile, 1149
Plutella xylostella, 1149
sex pheromone, 1149
trapping, 837, 1186, 1350
Cerambycidae, 837
detection, 1350
Drosophila suzukii, 1350
flight-intercept trap, 837
invasive fruit fly, 1186
male lure, 1186
monitoring, 837, 1350
surveillance, 1186
Tephritidae, 1186
wood-borer detection, 837
Tribolium castaneum, 670, 686, 1471
development, 1471
diatomaceous earth, 670
flour mill, 686
fumigation, 686
insecticidal potential, 670
integrated pest management (IPM), 686
natural zeolite, 670
nutritional quality, 1471
reproduction, 1471
Sitophilus oryzae, 670
spatial distribution, 686
stored product, 1471
Tribolium confusum,
Rhyzopertha dominica, 1088
short exposure, 1088
Sitophilus oryzae, 1088
spinetoram, 1088
Trichogramma achaeae,
cultivar, 2085
Nesidiocoris tenuis, 2085
release strategy, 2085
Tuta absoluta, 2085
Trichogramma chilonis,
insecticide, 1157
life-table parameter, 1157
sublethal effect, 1157
toxicity, 1157
Trichogramma japonicum,
biological control, 92
chemical control, 92
egg parasitoid, 92
Trichophysetis cretacea,
sex pheromone, 1566
(Z)-11-hexadecenal, 1566
(Z)-11-hexadecenol, 1566
(Z)-11-hexadecenyl acetate, 1566
Trifolium pratense,
Apion spp., 1620
biological control, 1620
Spintherus dubius, 1620
thiacloprid, 1620
trimethylamine,
ammonium acetate, 471
Ceratitis capitata, 471
fruit fly management, 471
spinosad, 471
triose phosphate isomerase,
cytochrome oxidase I, 418
haplotype, 418
Spodoptera frugiperda, 418
Triticum aestivum,
gall midge, 1988
Mayetiola destructor, 1988
seasonal abundance, 1988
sex pheromone, 1988
tritrophic interaction,
entomopathogenic nematode, 994
genetically modified crop, 994
meta-analysis, 994
Pectinophora gossypiella, 994
tritrophic interactions,
biological control, 161
ectoparasitoid, 161
leafminer, 161
sex ratio, 161
turfgrass, 1573, 2045
degree-day, 1573
developmental rate, 1573
Herpetogramma phaeopteralis, 1573
hunting billbug, 2045
nonlinear model, 1573
Sphenophorus venatus vestitus, 2045
Tuta absoluta,
cultivar, 2085
Nesidiocoris tenuis, 2085
release strategy, 2085
Trichogramma achaeae, 2085
two-sex life table,
age-stage, 1581
Aulacaspis yasumatsui, 1581
Cycas revoluta, 1581
Cycas taitungensis, 1581

U

ultrasound,
bed bug, 2107
Cimex lectularius, 2107
management, 2107
repellency, 2107
2-(undecyloxy)-ethanol,
aggregation pheromone, 2029
semiochemical, 2029
urban forest management,
economic analysis, 196
emerald ash borer, 196
preemptive removal, 196
urban insect pest,
spatial analysis, 1659
urban landscape, 1659
urban landscape,
spatial analysis, 1659
urban insect pest, 1659

V

Vaccinium corymbosum,
cryptic species, 1205
Dasineura oxycoccana, 1205
phenology, 1205
Vaccinium macrocarpon, 1205
Vaccinium corymbosum × *Vaccinium darrowi*,
economic injury level, 480
flower thrips, 480
Frankliniella bispinosa, 480
southern highbush blueberry, 480
Vaccinium corymbosum × *Vaccinium darrowi*, 480

Vaccinium macrocarpon,
cryptic species, 1205
Dasineura oxycoccana, 1205
phenology, 1205
Vaccinium corymbosum, 1205
varnished wood,
concrete, 718
methoprene, 718
residual efficacy, 718
surface treatment, 718
Varroa destructor, 313, 1895
Apis mellifera, 313, 1895
crop pollination, 313
genetic resistance, 313
migratory beekeeping, 313
miticide, 1895
pesticide, 1895
queen, 1895
vascular histochemistry,
oviposition puncture to larvae ratio, 598
plant resistance, 598
soybean petiole morphology, 598
soybean plant introduction, 598
vector, 1447, 1909
aphid, 1909
Begomovirus, 1447
DNA barcoding, 1909
host plant resistance, 1447
potato virus Y, 1909
transmission, 1447
vegetative insecticidal protein,
development, 964
eclosion, 964
hatching, 964
mating, 964
venom alkaloids,
Plutella xylostella, 1591
Solenopsis geminata, 1591
Solenopsis invicta, 1591
verbenone,
acetophenone, 1668
nonhost angiosperm volatile, 1668
pest management, 1668
Pinus ponderosa, 1668
verde plant bug,
cotton fleahopper, 896
Creontiades signatus, 896
pest sampling, 896
Pseudatomoscelis seriatus, 896
vineyard, 176, 1293
Empoasca vitis, 176
irrigation system, 176
leafhopper, 1293
regulated deficit irrigation (RDI), 1293
water, 1293
water stress, 176
virulence, 232, 344
baculovirus, 344
biotype, 232
DNA methylation, 232
geographic virus isolate, 344
Lymantria dispar asiatica, 344
pathogenicity, 344
salivary gland, 232
sodium bisulfite, 232
volatile,
(E)-β-caryophyllene, 120
fall armyworm, 120
maize, 120
resistance, 120

W

wash-off,
Popillia japonica, 1688
precipitation, 1688
water,
leafhopper, 1293
regulated deficit irrigation (RDI), 1293
vineyard, 1293
water stress,
Empoasca vitis, 176
irrigation system, 176
vineyard, 176
weed,
phloem-limited bacteria, 329
plant disease, 329
sap-sucking insect, 329
weevil,
kairomone, 169
Otiorhynchus, 169
plant odor, 169
strawberry, 169
weight study,
baseline data, 1024
basic model, 1024
50% lethal dose, 1024
proportionality, 1024
West Indian fruit fly,
Anastrepha obliqua, 2147
COI, 2147
mitochondrial DNA, 2147
sequencing, 2147
western flower thrip,
acrinathrin, 1006
cytochrome P450, 1006
insecticide resistance, 1006
over-expression, 1006
western tarnished plant bug,
biophysical model, 860
degree-day model, 860
temperature-dependent development, 860
wheat, 642, 854, 1057, 2180
adaptation, 1057
antibiosis, 2180
antixenosis, 2180
community index, 854
Diuraphis noxia, 1057
genetic diversity, 1057
Hessian fly, 642
intercropping, 854
natural enemy, 854
plant resistance, 1057
resistance, 642
Sitobion avenae, 854
susceptibility, 642
systemic induced susceptibility, 642
tolerance, 2180
wheat curl mite, 2180
wheat curl mite, 1801, 2180
Aceria tosichella, 1801
Aceria tulipae, 1801
antibiosis, 2180
antixenosis, 2180
tolerance, 2180
wheat, 2180
wheat streak mosaic, 1801
wheat line,
bird cherry-oat aphid, 592
plant resistance categories, 592
wheat streak mosaic,
Aceria tosichella, 1801
Aceria tulipae, 1801
wheat curl mite, 1801
white grub,
Cylas formicarius, 651
Diabrotica, 651
Ipomoea, 651
plant resistance, 651
whitefly, 48, 1401
acetamiprid, 1401
fitness cost, 1401
mitochondrial cytochrome oxidase I, 48
phylogenetic analysis, 48
resistance, 1401
white-tailed deer,
amitraz, 2207
collar, 2207
neckband, 2207
tick control, 2207
wood borer,
Asian longhorned beetle, 1645
bark beetle, 1645
emerald ash borer, 1645
exotic insect, 1645
wood-borer detection,
Cerambycidae, 837
flight-intercept trap, 837
monitoring, 837
trapping, 837
wood-boring insect,
Cerambycidae, 395
Fluon, 395
pheromone trap, 395

X

x-ray radiation,
  invasive species, 1971
  Lepidoptera, 1971
  phytosanitary treatment, 1971
  regulatory pest, 1971

Y

yellowmargined leaf beetle,
  crucifer, 1719
  field predation, 1719
  grower recommendation, 1719
  spined soldier bug, 1719

Z

zebra chip,
  areawide pest management, 1268
  Bactericera cockerelli, 1268
  insect vector, 1268
  Liberibacter, 1268

(Z)-11-hexadecenal,
  (Z)-11-hexadecenol, 1566
  (Z)-11-hexadecenyl acetate, 1566
  sex pheromone, 1566
  Trichophysetis cretacea, 1566
(3Z)-hexenol,
  Agrilus planipennis, 429
  dark green trap, 429
  manuka oil, 429
  phoebe oil, 429